

## ACTEMRA SC (S)

### Products Affected

- Actemra INJ 162MG/0.9ML
- Actemra Actpen

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: a) Either a trial and failure, contraindication, or intolerance (TF/C/I) to two of the following: Enbrel (etanercept), Humira (adalimumab), Rinvoq (upadacitinib), Xeljanz/Xeljanz XR (tofacitinib) or attestation demonstrating a trial may be inappropriate, OR b) For continuation of prior therapy. Giant Cell Arteritis (GCA) (Initial): Diagnosis of GCA. Trial and failure, contraindication, or intolerance to a glucocorticoid (eg, prednisone). Systemic Juvenile Idiopathic Arthritis (SJIA) (Initial): Diagnosis of active SJIA. Trial and failure, contraindication, or intolerance to one of the following: NSAID (eg, ibuprofen, naproxen), methotrexate, or systemic glucocorticoid (eg, prednisone). Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Diagnosis of active PJIA. One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior therapy. Systemic sclerosis-associated interstitial lung disease (SSc-ILD) (Initial): Diagnosis of SSc-ILD as documented by the following: a) Exclusion of other known causes of ILD AND b) One of the following: i) In patients not subjected to surgical lung biopsy, the presence of idiopathic interstitial pneumonia (eg, fibrotic nonspecific interstitial pneumonia [NSIP], usual interstitial pneumonia [UIP] and centrilobular fibrosis) pattern on high-resolution computed tomography (HRCT) revealing SSc-ILD or probable SSc-ILD, OR ii) In patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern revealing SSc-ILD or probable SSc-ILD.
Age Restrictions	N/A

<b>Prescriber Restrictions</b>	RA, GC, SJIA, PJIA (initial): Prescribed by or in consultation with a rheumatologist. SSc-ILD (initial): Prescribed by or in consultation with a pulmonologist or rheumatologist.
<b>Coverage Duration</b>	RA, GC, SJIA, PJIA, SSc-ILD (initial, reauth): 12 months
<b>Other Criteria</b>	RA, GC, SJIA, PJIA, SSc-ILD (Reauth): Documentation of positive clinical response to therapy.

## ACTIMMUNE (S)

---

### Products Affected

- Actimmune

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of one of the following: 1) Chronic granulomatous disease (CGD), or 2) severe malignant osteopetrosis (SMO).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## ADCIRCA (S)

### Products Affected

- Alyq
- Tadalafil TABS 20MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
Age Restrictions	N/A
Prescriber Restrictions	PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
Coverage Duration	PAH: Initial: 6 months. Reauth: 12 months.
Other Criteria	PAH (Reauth): Documentation of positive clinical response to therapy.

## ADEMPAS (S)

### Products Affected

- Adempas

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH AND PAH is symptomatic AND One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. Chronic thromboembolic pulmonary hypertension (CTEPH) (Initial): One of the following: A) Both of the following: 1) Diagnosis of inoperable or persistent/recurrent CTEPH and 2) CTEPH is symptomatic OR B) Patient is currently on any therapy for the diagnosis of CTEPH.
Age Restrictions	N/A
Prescriber Restrictions	PAH, CTEPH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
Coverage Duration	PAH, CTEPH: Initial: 6 months. Reauth: 12 months.
Other Criteria	PAH, CTEPH (Reauth): Documentation of positive clinical response to therapy.

## AFINITOR (S)

### Products Affected

- Afinitor TABS 10MG
- Everolimus TABS 10MG, 2.5MG, 5MG, 7.5MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Subependymal Giant Cell Astrocytoma (SEGA) associated with tuberous sclerosis complex (TSC): Diagnosis of SEGA associated with TSC that requires therapeutic intervention but patient is not a candidate for curative surgical resection. Renal cell carcinoma: Diagnosis of advanced or metastatic renal cell carcinoma AND trial and failure, contraindication, or intolerance to SUTENT (sunitinib) or NEXAVAR (sorafenib). Neuroendocrine tumors of pancreatic origin (pNET): Diagnosis of progressive pNET that are unresectable, locally advanced, or metastatic. Renal angiomyolipoma: Diagnosis of renal angiomyolipoma and TSC AND Patient does not require immediate surgery. Breast Cancer: Patient is a postmenopausal woman AND Diagnosis of advanced hormone receptor-positive, HER2-negative breast cancer AND trial and failure, contraindication, or intolerance to FEMARA (letrozole) or ARIMIDEX (anastrozole) AND used in combination with AROMASIN (exemestane). Neuroendocrine tumors of gastrointestinal (GI) or lung origin: Diagnosis of progressive, well-differentiated, non-functional NET of GI or lung origin AND patient has unresectable, locally advanced or metastatic disease.
Age Restrictions	N/A
Prescriber Restrictions	All uses: Prescribed by or in consultation with an oncologist
Coverage Duration	All uses: 12 months
Other Criteria	All Indications: Approve for continuation of prior therapy.

## AFINITOR DISPERZ (S)

### Products Affected

- Afinitor Disperz
- Everolimus TBSO

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Subependymal Giant Cell Astrocytoma (SEGA): Diagnosis of SEGA associated with tuberous sclerosis complex (TSC) that requires therapeutic intervention but patient is not a candidate for curative surgical resection. TSC-associated partial-onset seizures: Diagnosis of TSC-associated partial-onset seizures. Used as adjunctive therapy.
Age Restrictions	N/A
Prescriber Restrictions	SEGA: Prescribed by or in consultation with an oncologist. TSC-associated partial-onset seizures: Prescribed by or in consultation with a neurologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# AIMOVIG (S)

## Products Affected

- Aimovig

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Episodic Migraines (EM) (initial): Diagnosis of EM. Patient has 4 to 14 migraine days per month, but no more than 14 headache days per month.</p> <p>Chronic Migraines (CM) (initial): Diagnosis of CM. Medication overuse headache has been considered and potentially offending medication(s) have been discontinued. Patient has greater than or equal to 15 headache days per month, of which at least 8 must be migraine days for at least 3 months.</p> <p>All Indications (initial): Two of the following: a) History of failure (after at least a two month trial) or intolerance to Elavil (amitriptyline) or Effexor (venlafaxine), OR patient has a contraindication to both Elavil (amitriptyline) and Effexor (venlafaxine), b) History of failure (after at least a two month trial) or intolerance to Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate), OR patient has a contraindication to both Depakote/Depakote ER (divalproex sodium) and Topamax (topiramate), c) History of failure (after at least a two month trial) or intolerance to one of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol, OR patient has a contraindication to all of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol, or d) History of failure (after at least a two month trial) or intolerance to Atacand (candesartan), OR patient has a contraindication to Atacand (candesartan). Medication will not be used in combination with another injectable CGRP inhibitor.</p>
Age Restrictions	EM, CM (initial): 18 years of age or older.
Prescriber Restrictions	EM, CM (initial, reauth): Prescribed by or in consultation with a neurologist, headache specialist, or pain specialist.
Coverage Duration	EM, CM (initial): 6 months. EM, CM (reauth): 12 months.



<b>Other Criteria</b>	<p>EM, CM (reauth): Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity. Use of acute migraine medications (e.g., non-steroidal anti-inflammatory drugs [NSAIDs] [e.g., ibuprofen, naproxen], triptans [e.g., eletriptan, rizatriptan, sumatriptan]) has decreased since the start of CGRP therapy. Medication will not be used in combination with another injectable CGRP inhibitor. CM (reauth): Patient continues to be monitored for medication overuse headache.</p>
-----------------------	---

## ALDURAZYME (S)

---

### Products Affected

- Aldurazyme

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Mucopolysaccharidosis I: confirmed diagnosis of Hurler and Hurler-Scheie forms of Mucopolysaccharidosis I (MPS I), OR confirmed diagnosis of Scheie form of Mucopolysaccharidosis I (MPS I) who have moderate to severe symptoms.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

## ALECENSA (S)

---

### Products Affected

- Alecensa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): A) Diagnosis of metastatic NSCLC AND B) Patient has anaplastic lymphoma kinase (ALK)-positive disease as detected with a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## ALPHA-1 PROTEINASE INHIBITOR, NON-PREFERRED (S)

### Products Affected

- Zemaira
- Aralast Np INJ 1000MG, 500MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Alpha-1 antitrypsin (AAT) deficiency: Diagnosis of congenital AAT deficiency. Diagnosis of emphysema. Continued optimal conventional treatment for emphysema (e.g., bronchodilators). One of the following: 1) PiZZ, PiZ(null), or Pi(null)(null) protein phenotypes (homozygous) OR 2) other rare AAT disease genotypes associated with pre-treatment serum AAT level less than 11 µM/L [e.g., Pi(Malton, Malton), Pi(SZ)]. One of the following: Circulating pre-treatment serum AAT level less than 11 µM/L (which corresponds to less than 80 mg/dL if measured by radial immunodiffusion or less than 57 mg/dL if measured by nephelometry) OR the patient has a concomitant diagnosis of necrotizing panniculitis. Trial and failure, or intolerance to Prolastin.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

# ALPHA-1 PROTEINASE INHIBITOR, PROLASTIN (S)

## Products Affected

- Prolastin-c INJ 1000MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Alpha-1 antitrypsin (AAT) deficiency: Diagnosis of congenital AAT deficiency. Diagnosis of emphysema. Continued optimal conventional treatment for emphysema (e.g., bronchodilators). One of the following: 1) PiZZ, PiZ(null), or Pi(null)(null) protein phenotypes (homozygous) OR 2) other rare AAT disease genotypes associated with pre-treatment serum AAT level less than 11 µM/L [e.g., Pi(Malton, Malton), Pi(SZ)]. Circulating pre-treatment serum AAT level less than 11 µM/L (which corresponds to less than 80 mg/dL if measured by radial immunodiffusion or less than 57 mg/dL if measured by nephelometry), unless the patient has a concomitant diagnosis of necrotizing panniculitis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

## ALUNBRIG (S)

---

### Products Affected

- Alunbrig

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC. Patient has an anaplastic lymphoma kinase (ALK)-positive tumor as detected with a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## AMPYRA (S)

### Products Affected

- Dalfampridine Er

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple Sclerosis (MS) (initial): Diagnosis of MS. Physician confirmation that patient has difficulty walking (eg, timed 25 foot walk test). One of the following: expanded disability status scale (EDSS) score less than or equal to 7, or not restricted to using a wheelchair (if EDSS is not measured).
Age Restrictions	N/A
Prescriber Restrictions	MS (initial): Prescribed by or in consultation with a neurologist.
Coverage Duration	MS (Initial): 6 months. (Reauth): 12 months.
Other Criteria	MS (Reauth): Physician confirmation that the patient's walking improved with therapy. One of the following: EDSS score less than or equal to 7, or not restricted to using a wheelchair (if EDSS is not measured).

# ARCALYST (S)

## Products Affected

- Arcalyst

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cryopyrin-Associated Periodic Syndromes (CAPS) (Initial): Diagnosis of CAPS, including Familial Cold Auto-inflammatory Syndrome (FCAS) and/or Muckle-Wells Syndrome (MWS). The medication will not be used in combination with another biologic. Deficiency of Interleukin-1 Receptor Antagonist (DIRA): Diagnosis of DIRA. Patient weighs at least 10 kg. Patient is currently in remission (e.g., no fever, skin rash, and bone pain/no radiological evidence of active bone lesions/C-reactive protein [CRP] less than 5 mg/L). Recurrent Pericarditis (Initial): Diagnosis of recurrent pericarditis as evidenced by at least 2 episodes that occur a minimum of 4 to 6 weeks apart. Trial and failure, contraindication, or intolerance (TF/C/I) to at least one of the following: nonsteroidal anti-inflammatory drugs (e.g., ibuprofen, naproxen), colchicine, or corticosteroids (e.g., prednisone).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CAPS (initial): Prescribed by or in consultation with an immunologist, allergist, dermatologist, rheumatologist, neurologist or other medical specialist. Recurrent Pericarditis (initial): Prescribed by or in consultation with a cardiologist.
<b>Coverage Duration</b>	CAPS, Recurrent Pericarditis (initial, reauth): 12 months. DIRA: 12 months.
<b>Other Criteria</b>	CAPS (Reauth): Patient has experienced disease stability or improvement in clinical symptoms while on therapy as evidence by one of the following: A) improvement in rash, fever, joint pain, headache, conjunctivitis, B) decreased number of disease flare days, C) normalization of inflammatory markers (CRP, ESR, SAA), D) corticosteroid dose reduction, OR E) improvement in MD global score or active joint count. Recurrent Pericarditis (Reauth): Documentation of positive clinical response to therapy.



## AURYXIA (S)

---

### Products Affected

- Auryxia

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Exclude if used for iron deficiency anemia in chronic kidney disease (CKD) not on dialysis.
Required Medical Information	Hyperphosphatemia in chronic kidney disease: Diagnosis of hyperphosphatemia. Patient has chronic kidney disease (CKD). Patient is on dialysis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

# AUSTEDO (S)

## Products Affected

- Austedo

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chorea associated with Huntington's disease (initial): Diagnosis of Chorea associated with Huntington's disease. Tardive dyskinesia (initial): Diagnosis of moderate to severe tardive dyskinesia. One of the following: 1) Patient has persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering, or discontinuation of the offending medication or 2) Patient is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Huntington's disease chorea (initial): Prescribed by a neurologist. Tardive dyskinesia (initial): Prescribed by or in consultation with a neurologist or psychiatrist.
<b>Coverage Duration</b>	Initial: 3 months. Reauth: 12 months
<b>Other Criteria</b>	All indications (Reauth): Documentation of positive clinical response to therapy.

## AYVAKIT (S)

### Products Affected

- Ayvakit

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Gastrointestinal stromal tumor (GIST): Diagnosis of GIST. Disease is one of the following: unresectable or metastatic. Presence of platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations. Advanced Systemic Mastocytosis (AdvSM): Diagnosis of AdvSM. Patient has one of the following: a) aggressive systemic mastocytosis (ASM), b) systemic mastocytosis with an associated hematological neoplasm (SM-AHN), or c) mast cell leukemia (MCL).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	GIST: Prescribed by or in consultation with an oncologist. AdvSM: Prescribed by or in consultation with an oncologist/hematologist, allergist, or immunologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## BAFIERTAM (S)

### Products Affected

- Bafiertam

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple Sclerosis (MS) (initial): Diagnosis of a relapsing form of MS (e.g., clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions). One of the following: a) Failure after a trial of at least 4 weeks, contraindication, or intolerance to two of the following disease-modifying therapies for MS: 1) Aubagio (teriflunomide), 2) Gilenya (fingolimod), or 3) Brand Tecfidera/generic dimethyl fumarate, OR b) for continuation of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	MS (initial, reauth): Prescribed by or in consultation with a neurologist
Coverage Duration	MS (initial, reauth): 12 months
Other Criteria	MS (reauth): Documentation of positive clinical response to therapy (e.g., stability in radiologic disease activity, clinical relapses, disease progression).

## BALVERSA (S)

### Products Affected

- Balversa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Urothelial Carcinoma: Diagnosis of urothelial carcinoma (UC). One of the following: Locally advanced or Metastatic AND Patient has fibroblast growth factor receptor (FGFR) 3 or FGFR2 genetic alterations as detected by an U.S. Food and Drug Administration (FDA)-approved test (therascreen FGFR RGQ RT-PCR Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). One of the following: 1) Patient has progressed during or following at least one line of prior chemotherapy or immunotherapy OR 2) Patient has progressed within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# BENLYSTA (S)

## Products Affected

- Benlysta

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Systemic lupus erythematosus (SLE) (init): Diagnosis of active SLE. Autoantibody positive (ie, anti-nuclear antibody [ANA] titer greater than or equal to 1:80 or anti-dsDNA level greater than or equal to 30 IU/mL). Currently receiving at least one standard of care treatment for active SLE (eg, antimalarials [eg, Plaquenil (hydroxychloroquine)], corticosteroids [eg, prednisone], or immunosuppressants [eg, methotrexate, Imuran (azathioprine)]). Lupus Nephritis (init): Diagnosis of active lupus nephritis. Currently receiving standard of care treatment for active lupus nephritis (e.g., corticosteroids [e.g., prednisone] with mycophenolate or cyclophosphamide).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	SLE (init): Prescribed by or in consultation with a rheumatologist. Lupus Nephritis (init): Prescribed by or in consultation with a nephrologist or rheumatologist.
<b>Coverage Duration</b>	SLE, Lupus Nephritis (init, reauth): 6 months
<b>Other Criteria</b>	SLE, Lupus Nephritis (reauth): Documentation of positive clinical response to therapy.

## BESREMI (S)

### Products Affected

- Besremi

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of polycythemia vera as confirmed by all of the following: 1) One of the following: a) Hemoglobin greater than 16.5 g/dL for men or hemoglobin greater than 16.0 g/dL for women, b) Hematocrit greater than 49% for men or hematocrit greater than 48% for women, or c) Increased red cell mass, AND 2) Bone marrow biopsy showing hypercellularity for age with trilineage growth (panmyelosis) including prominent erythroid, granulocytic and megakaryocytic proliferation with pleomorphic, mature megakaryocytes, AND 3) One of the following: a) Presence of JAK2 or JAK2 exon 12 mutation or b) Subnormal serum erythropoietin level. Both of the following: 1) Trial and failure, contraindication or intolerance (TF/C/I) to hydroxyurea, AND 2) TF/C/I to one interferon therapy (e.g., Intron A, Pegasys, etc).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## BOSULIF (S)

---

### Products Affected

- Bosulif

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic myelogenous/myeloid leukemia (CML): Diagnosis of Philadelphia chromosome-positive (Ph+) CML.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.



# BRAFTOVI (S)

## Products Affected

- Braftovi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Melanoma: Diagnosis of unresectable melanoma or metastatic melanoma. Cancer is BRAF V600E or V600K mutant type (MT) as detected by a U.S. Food and Drug Administration (FDA)-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Used in combination with Mektovi (binimetinib). Colorectal Cancer: One of the following diagnoses: Colon Cancer or Rectal Cancer. One of the following: 1) Unresectable or advanced disease or 2) Metastatic disease. Patient has received prior therapy. Cancer is BRAF V600E mutant type as detected by a U.S. Food and Drug Administration (FDA)-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Used in combination with Erbitux (cetuximab).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy

## BRIVIACT (S)

---

### Products Affected

- Briviact SOLN
- Briviact TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Partial-onset seizures: Diagnosis of partial-onset seizures.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## BRUKINSA (S)

### Products Affected

- Brukinsa

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Mantle Cell Lymphoma (MCL): Diagnosis of relapsed or refractory MCL. Trial and failure, contraindication, or intolerance to at least ONE combination treatment of rituximab and chemotherapy (e.g., BR, R-CHOP, R-CVP, R-FCM). Waldenstrom's Macroglobulinemia (WM)/Lymphoplasmacytic Lymphoma (LPL): Diagnosis of WM/LPL. Marginal Zone Lymphoma (MZL): Diagnosis of MZL. Disease is relapsed or refractory. Patient has received at least one prior anti-CD20-based regimen for MZL (e.g., rituximab, obinutuzumab).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist/hematologist.
<b>Coverage Duration</b>	12 months.
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## CABLIVI (S)

### Products Affected

- Cablivi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acquired thrombotic thrombocytopenic purpura (aTTP): Diagnosis of aTTP. First dose was/will be administered by a healthcare provider as a bolus intravenous injection. Used in combination with immunosuppressive therapy (e.g. rituximab, glucocorticoids). One of the following: 1) Used in combination with plasma exchange or 2) both of the following: patient has completed plasma exchange and less than 59 days have or will have elapsed beyond the last plasma exchange.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist.
Coverage Duration	3 months
Other Criteria	N/A

# CABOMETYX (S)

## Products Affected

- Cabometyx

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Renal cell carcinoma (RCC): Diagnosis of RCC. RCC is advanced. Hepatocellular Carcinoma (HCC): Diagnosis of HCC. One of the following: a) Trial and failure, contraindication, or intolerance to Nexavar (sorafenib tosylate), or b) Patient has metastatic disease, or c) Patient has extensive liver tumor burden, or d) Patient is inoperable by performance status or comorbidity (local disease or local disease with minimal extrahepatic disease only), or e) Disease is unresectable. Differentiated Thyroid Cancer (DTC): Diagnosis of DTC. Disease is one of the following: a) locally advanced or b) metastatic. Disease has progressed following prior VEGFR-targeted therapy (e.g., Lenvima [lenvatinib], Nexavar [sorafenib]). Disease or patient is refractory to radioactive iodine treatment or ineligible.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RCC: Prescribed by or in consultation with one of the following: an oncologist or nephrologist. HCC: Prescribed by or in consultation with one of the following: oncologist, hepatologist, or gastroenterologist. DTC: Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## CALQUENCE (S)

---

### Products Affected

- Calquence

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Mantle Cell Lymphoma: Diagnosis of mantle cell lymphoma (MCL) AND patient has received at least one prior therapy for MCL. Chronic Lymphocytic Leukemia (CLL) or Small Lymphocytic Lymphoma (SLL): Diagnosis of CLL or SLL.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## CAPRELSA (S)

---

### Products Affected

- Caprelsa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Thyroid Cancer: Diagnosis of medullary thyroid cancer (MTC).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with oncologist or endocrinologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## CAYSTON (S)

---

### Products Affected

- Cayston

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cystic fibrosis (CF) (Initial, Reauth): Diagnosis of CF AND Patient has evidence of Pseudomonas aeruginosa in the lungs.
Age Restrictions	CF (Initial): 7 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	CF (Initial, reauth): 12 months
Other Criteria	CF (Reauth): Patient is benefiting from treatment (i.e. improvement in lung function [forced expiratory volume in one second (FEV1)], decreased number of pulmonary exacerbations).



# CERDELGA (S)

## Products Affected

- Cerdelga

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Gaucher disease (Initial): Diagnosis of Gaucher disease type 1. Patient is an extensive metabolizer (EM), intermediate metabolizer (IM), or poor metabolizer (PM) of cytochrome P450 enzyme (CYP) 2D6 as detected by an FDA-cleared test.
Age Restrictions	Gaucher disease (initial): 18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Gaucher disease (initial, reauth): 12 months
Other Criteria	Gaucher disease (Reauth): Patient's condition has not progressed, as defined by ALL of the following: A) Hemoglobin level decreased greater than 1.5 g/dL from baseline, AND B) Platelet count decreased greater than 25% from baseline, AND C) Spleen volume increased greater than 25% from baseline, AND D) Liver volume increased greater than 20% from baseline.

# CHOLBAM (S)

## Products Affected

- Cholbam

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Bile acid synthesis disorders due to single enzyme defects (BAS) (initial): diagnosis of a bile acid synthesis disorder due to a single enzyme defect. Peroxisomal disorders (PD) (initial): All of the following: 1) diagnosis of peroxisomal disorder, 2) patient exhibits at least one of the following: a) liver disease (eg, jaundice, elevated serum transaminases), OR b) steatorrhea, OR c) complications from decreased fat-soluble vitamin absorption (eg, poor growth), AND 3) will be used as an adjunctive treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses (initial): Prescribed by a hepatologist, medical geneticist, pediatric gastroenterologist, OR other specialist that treats inborn errors of metabolism.
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	All uses (reauth): documentation of positive clinical response to therapy.

## CICLOPIROX (S)

### Products Affected

- Ciclodan SOLN
- Ciclopirox Nail Lacquer

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	All of the following: 1) Patient does not have lunula (matrix) involvement, 2) one of the following: a) Diagnosis of onychomycosis of the toenails, OR b) Diagnosis of onychomycosis of the fingernails, 3) Diagnosis of onychomycosis has been confirmed by one of the following: a) positive potassium hydroxide (KOH) preparation, OR b) culture, OR c) histology, 4) If toenail onychomycosis, patient has mild to moderate disease involving at least 1 target toenail, AND 5) Trial and failure, contraindication, or intolerance to oral terbinafine.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	48 weeks.
Other Criteria	N/A

## CINRYZE (S)

### Products Affected

- Cinryze

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prophylaxis of hereditary angioedema (HAE) attacks: Diagnosis of HAE. Diagnosis has been confirmed by C1 inhibitor (C1-INh) deficiency or dysfunction (Type I or II HAE) as documented by one of the following: a) C1-INH antigenic level below the lower limit of normal OR b) C1-INH functional level below the lower limit of normal. For prophylaxis against HAE attacks.
Age Restrictions	N/A
Prescriber Restrictions	HAE (prophylaxis): Prescribed by or in consultation with an immunologist or an allergist
Coverage Duration	12 months
Other Criteria	N/A

## COMETRIQ (S)

---

### Products Affected

- Cometriq

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Medullary thyroid cancer (MTC): Diagnosis of Metastatic MTC.
Age Restrictions	N/A
Prescriber Restrictions	MTC: Prescribed by or in consultation with an oncologist/hematologist or endocrinologist.
Coverage Duration	All uses: 12 months
Other Criteria	Approve for continuation of prior therapy.

## COPIKTRA (S)

### Products Affected

- Copiktra

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): Diagnosis of CLL or SLL. Disease is relapsed or refractory. Trial and failure, contraindication, or intolerance to at least two prior therapies for CLL/SLL (e.g., Leukeran [chlorambucil], Gazyva [obinutuzumab], Arzerra [ofatumumab], Bendeka [bendamustine], Imbruvica [ibrutinib], Rituxan [rituximab], etc.). Follicular Lymphoma: Diagnosis of follicular lymphoma. Disease is relapsed or refractory. Trial and failure, contraindication, or intolerance to at least two prior systemic therapies for follicular lymphoma (e.g., Leukeran [chlorambucil], Gazyva [obinutuzumab], Arzerra [ofatumumab], Bendeka [bendamustine], Imbruvica [ibrutinib], Rituxan [rituximab], etc.).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

# CORLANOR (S)

## Products Affected

- Corlanor TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic heart failure (CHF) (initial): Diagnosis of CHF. Patient has NYHA Class II, III, or IV symptoms. Patient has a left ventricular ejection fraction less than or equal to 35%. Patient is in sinus rhythm. Patient has a resting heart rate of greater than or equal to 70 beats per minute. One of the following: patient is on a beta-blocker (e.g., bisoprolol, carvedilol, metoprolol succinate extended release) at a maximally tolerated dose, or patient has a contraindication or intolerance to beta-blocker therapy. Patient has been hospitalized for worsening HF in the previous 12 months. Trial and failure, contraindication, or intolerance to maximally tolerated doses of an ACE inhibitor (e.g., captopril, enalapril, lisinopril) or ARB (e.g., candesartan, losartan, valsartan). Dilated Cardiomyopathy (DCM) (initial): Diagnosis of heart failure due to DCM. Patient has NYHA Class II, III, or, IV symptoms. Patient is in sinus rhythm. Patient has an elevated heart rate. Trial and failure, contraindication or intolerance to one of the following: 1) Beta blocker (e.g., bisoprolol, metoprolol succinate extended release), 2) Angiotensin-converting enzyme (ACE) inhibitor (e.g., captopril, enalapril), or 3) Diuretic Agent (e.g., spironolactone, furosemide).
Age Restrictions	N/A
Prescriber Restrictions	CHF, DCM (initial): Prescribed by or in consultation with a cardiologist
Coverage Duration	CHF, DCM (initial, reauth): 12 months
Other Criteria	CHF, DCM (reauth): Documentation of positive clinical response to therapy.

# COSENTYX (S)

## Products Affected

- Cosentyx

- Cosentyx Sensoready Pen

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. One of the following: Trial and failure, contraindication (eg, safety concerns, not indicated for patient's age/weight), or intolerance to one of the following: Enbrel (etanercept), Humira (adalimumab), Skyrizi (risankizumab), OR for continuation of prior therapy. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: a) Either trial and failure, contraindication (eg, safety concerns, not indicated for patient's age/weight), or intolerance to two of the following, or attestation demonstrating a trial may be inappropriate: Enbrel (etanercept), Humira (adalimumab), Skyrizi (risankizumab-rzaa), Rinvoq (upadacitinib), Xeljanz/XR (tofacitinib/ER), OR b) for continuation of prior therapy. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. One of the following: Trial and failure, contraindication, or intolerance to Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior therapy. Non-radiographic axial spondyloarthritis (nr-axSpA, initial): Dx of active nr-axSpA with objective signs of inflammation (eg, C-reactive protein [CRP] levels above the upper limit of normal and/or sacroiliitis on magnetic resonance imaging [MRI], indicative of inflammatory disease, but without definitive radiographic evidence of structural damage on sacroiliac joints.) TF/C/I to two non-steroidal anti-inflammatory drugs (NSAIDs) (eg, ibuprofen, meloxicam, naproxen). Enthesitis-Related Arthritis (ERA) (Initial): Diagnosis of active ERA. Trial and failure, contraindication, or intolerance to TWO non-steroidal anti-inflammatory drugs (NSAIDs) (eg, ibuprofen, meloxicam, naproxen).</p>
Age Restrictions	ERA (initial): Patient is 4 years of age or older.



<b>Prescriber Restrictions</b>	Plaque psoriasis (initial): Prescribed by or in consultation with a dermatologist. PsA (initial): Prescribed by or in consultation with a rheumatologist or dermatologist. AS, nr-axSpA, ERA (initial): Prescribed by or in consultation with a rheumatologist.
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	PsA, AS, nr-axSpA (Reauth): Documentation of positive clinical response to therapy. Psoriasis (Reauth): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline. ERA (Reauth): Documentation of a positive clinical response to therapy as evidenced by at least one of the following: Reduction in the total active (swollen and tender) joint count from baseline, OR improvement in symptoms (eg, pain, stiffness, inflammation) from baseline.

## COTELLIC (S)

---

### Products Affected

- Cotellic

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Melanoma: Diagnosis of unresectable or metastatic melanoma. Patient has a BRAF V600E or V600K mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., cobas 4800 BRAF V600 Mutation Test) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Used in combination with vemurafenib.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## CYSTARAN (S)

---

### Products Affected

- Cystaran

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cystinosis: Diagnosis of cystinosis, confirmed by elevated leukocyte cystine levels (LCL), genetic analysis of the CTNS gene or corneal cystine crystal accumulation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

## D.H.E. 45 (S)

### Products Affected

- Dihydroergotamine Mesylate INJ

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Migraines (initial): Diagnosis of migraine headaches with or without aura. Will be used for the acute treatment of migraine. One of the following: Trial and failure or intolerance to one triptan (e.g., eletriptan, rizatriptan, sumatriptan) or contraindication to all triptans. If patient has 4 or more headache days per month, patient must meet one of the following: a) currently being treated with Elavil (amitriptyline) or Effexor (venlafaxine) unless there is a contraindication or intolerance to these medications, OR b) currently being treated with Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) unless there is a contraindication or intolerance to these medications, OR c) currently being treated with a beta blocker (i.e., atenolol, propranolol, nadolol, timolol, or metoprolol) unless there is a contraindication or intolerance to these medications. Cluster Headaches (CH) (initial): Diagnosis of cluster headache. Trial and failure, contraindication, or intolerance to sumatriptan injection.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial, Reauth: Prescribed by or in consultation with a neurologist, headache specialist, or pain specialist.
<b>Coverage Duration</b>	Migraines, CH (initial): 3 months. Migraines, CH (reauth): 12 months.
<b>Other Criteria</b>	Migraines (reauth): Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea). CH (reauth): Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity.

## DALIRESP (S)

---

### Products Affected

- Daliresp

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic Obstructive Pulmonary Disease (COPD) (initial): Diagnosis of COPD. History of COPD exacerbations which required the use of systemic corticosteroids, antibiotics, or hospital admission. Trial and failure, intolerance, or contraindication to two prior therapies for COPD (e.g., Combivent, Spiriva).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	COPD (init, reauth): 12 months
Other Criteria	COPD (reauth): Documentation of positive clinical response to therapy.

## DANYELZA (S)

---

### Products Affected

- Danyelza

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Neuroblastoma: Diagnosis of high-risk neuroblastoma in bone or bone marrow. Disease is relapsed or refractory. Used in combination with granulocyte-macrophage colony-stimulating factor [e.g., Leukine (sargramostim)]. Patient has had prior therapy with one of the following responses: partial response, minor response, or stable disease.
Age Restrictions	Neuroblastoma: Patient is 1 year of age or older.
Prescriber Restrictions	Neuroblastoma: Prescribed by or in consultation with an oncologist or hematologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# DARAPRIM (S)

## Products Affected

- Pyrimethamine TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Toxoplasmosis: 1) Patient is using pyrimethamine for the treatment of toxoplasmic encephalitis, secondary prophylaxis of toxoplasmic encephalitis, or treatment of congenital toxoplasmosis OR 2) Patient is using pyrimethamine for the primary prophylaxis of toxoplasmic encephalitis, patient has experienced intolerance to prior prophylaxis with trimethoprim-sulfamethoxazole (TMP-SMX), and one of the following: patient has been re-challenged with TMP-SMX using a desensitization protocol and is still unable to tolerate, or evidence of life-threatening reaction to TMP-SMX in the past (eg, toxic epidermal necrolysis, Stevens-Johnson syndrome). Malaria: Patient is using pyrimethamine for the treatment of acute malaria or chemoprophylaxis of malaria. Patient does not have megaloblastic anemia due to folate deficiency. The provider acknowledges that pyrimethamine is not recommended by the Centers for Disease Control and Prevention (CDC) for the treatment and/or prophylaxis of malaria.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist
Coverage Duration	12 months
Other Criteria	Toxoplasmosis only: Approve for continuation of prior therapy.

# DARZALEX FASPRO (S)

## Products Affected

- Darzalex Faspro

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Relapsed/Refractory Multiple Myeloma (MM): Diagnosis of MM. One of the following: A) Both of the following: Used as monotherapy and One of the following: i) Patient has received at least three prior treatment regimens which included both a proteasome inhibitor (e.g., bortezomib [Velcade], carfilzomib [Kyprolis]) and an immunomodulatory agent (e.g., lenalidomide [Revlimid], thalidomide [Thalomid]) or ii) patient is double-refractory to a proteasome inhibitor and an immunomodulatory agent. OR B) Both of the following: used in combination with one of the following treatment regimens: lenalidomide and dexamethasone, or bortezomib and dexamethasone, AND patient has received at least one prior therapy (e.g., bortezomib [Velcade], carfilzomib [Kyprolis], ixazomib [Ninlaro], lenalidomide [Revlimid], thalidomide [Thalomid]). Newly Diagnosed MM: Newly diagnosed MM. One of the following: A) Both of the following: pPatient is ineligible for autologous stem cell transplant AND one of the following: 1) used in combination with all of the following: bortezomib, melphalan, and prednisone or 2) used in combination with both of the following: lenalidomide and dexamethasone. OR B) Both of the following: patient is eligible for autologous stem cell transplant AND used in combination with all of the following: bortezomib, thalidomide, and dexamethasone. Light Chain (AL) Amyloidosis: Newly diagnosed light chain (AL) amyloidosis. Used in combination with all of the following: bortezomib, cyclophosphamide, and dexamethasone. All of the following: patient does not have New York Association (NYHA) Class IIIB disease, patient does not have NYHA class IV disease, and patient does not have Mayo Stage IIIB disease.</p>
Age Restrictions	N/A
Prescriber Restrictions	<p>MM: Prescribed by or in consultation with an oncologist/hematologist. Light Chain (AL) Amyloidosis: Prescribed by or in consultation with a hematologist.</p>



<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy

## DAURISMO (S)

---

### Products Affected

- Daurismo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute myeloid leukemia (AML): Diagnosis of newly-diagnosed acute myeloid leukemia (AML) AND Used in combination with low-dose cytarabine AND One of the following: 1) Patient is greater than or equal to 75 years old, or 2) Patient has comorbidities that preclude the use of intensive induction chemotherapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# DEFERASIROX (S)

## Products Affected

- Deferasirox

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic Iron Overload Due to Blood Transfusions (Initial): Diagnosis of chronic iron overload due to blood transfusions (transfusional hemosiderosis). Patient has a baseline ferritin level more than 1,000 mcg/L. Patient has required the transfusion of at least 100 mL/kg packed red blood cells. Myelodysplastic Syndrome (MDS) (Initial): Diagnosis of MDS. Patient has Low or Intermediate-1 disease or is a potential transplant patient. Patient has received more than 20 red blood cell transfusions. Chronic iron overload due to non-transfusion-dependent thalassemia (NTDT) (Initial): Diagnosis of chronic iron overload due to NTDT. Liver iron concentration (LIC) 5 milligrams of iron per gram of liver dry weight (mg Fe/g dw) or higher. Serum ferritin level greater than 300 mcg/L.
<b>Age Restrictions</b>	Iron Overload Due to Blood Transfusions (initial): 2 years of age or older. NTDT (initial): 10 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Iron Overload Due to Blood Transfusions, MDS (initial, reauth):12 mo. NTDT (initial, reauth): 6mo.
<b>Other Criteria</b>	Iron Overload Due to Blood Transfusions, MDS (Reauth): Patient experienced a reduction from baseline in serum ferritin level or LIC. NTDT (Reauth): Patient has LIC 3 mg Fe/g dw or higher. Patient experienced a reduction from baseline in serum ferritin level or LIC.

## DIACOMIT (S)

---

### Products Affected

- Diacomit

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of seizures associated with Dravet syndrome (DS). Used in combination with clobazam.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# DUPIXENT (S)

## Products Affected

- Dupixent

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Atopic dermatitis (AD) (init): Diagnosis (dx) of mod to severe AD. Trial and failure, contraindication, or intolerance (TF/C/I) to one med to high potency topical corticosteroid (eg, betamethasone, triamcinolone). One of the following: A) TF/C/I to pimecrolimus topical cream, unless patient is not a candidate for pimecrolimus therapy (e.g., immunocompromised, severe atopic dermatitis), B) TF/C/I to tacrolimus topical ointment, unless patient is not a candidate for tacrolimus ointment therapy (e.g., immunocompromised), or C) Eucrisa (crisaborole). Eosinophilic Asthma (EA) (init): Dx of mod to severe asthma. Asthma is an eosinophilic phenotype as defined by a baseline (pre-treatment) peripheral blood eosinophil level greater than or equal to 150 cells/microliter. One of the following: 1) Patient has had at least one or more asthma exacerbations requiring systemic corticosteroids (eg, prednisone) within the past 12 mo, 2) Any prior intubation for an asthma exacerbation, or 3) Prior asthma-related hospitalization within the past 12 mo. One of the following: a) TF/C/I to Fasenra (benralizumab), Nucala (mepolizumab), or Cinqair (reslizumab) or b) For continuation of prior therapy. Corticosteroid Dependent Asthma (CDA) (init): Dx of mod to severe asthma. Patient is currently dependent on oral corticosteroids for the treatment of asthma. EA, CDA (init): Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) High-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist (eg, montelukast), long-acting beta-2 agonist (LABA) (eg, salmeterol), tiotropium], OR b) One max-dosed combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone/vilanterol)].</p>
Age Restrictions	Asthma, Atopic dermatitis (initial): Age greater than or equal to 6 years. CRSwNP: no age restriction.

<b>Prescriber Restrictions</b>	Atopic dermatitis (Initial): Prescribed by or in consultation with one of the following: dermatologist, allergist/immunologist. Asthma (initial, reauth): Prescribed by or in consultation with a pulmonologist or allergist/immunologist. CRSwNP (initial, reauth): Prescribed by or in consultation with an otolaryngologist, allergist/immunologist, or pulmonologist.
<b>Coverage Duration</b>	Atopic Dermatitis, CRSwNP (Init/Reauth): 12 months. Asthma (Init): 6 mo. Asthma (reauth): 12 mo.
<b>Other Criteria</b>	Chronic rhinosinusitis with nasal polyposis (CRSwNP) (initial): Diagnosis of CRSwNP. Unless contraindicated, the patient has had an inadequate response to 2 months of treatment with an intranasal corticosteroid (eg, fluticasone, mometasone). Used in combination with another agent for CRSwNP. Atopic dermatitis (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in body surface area involvement, reduction in pruritus severity). EA (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in exacerbations, improvement in forced expiratory volume in 1 second [FEV1], decreased use of rescue medications). CDA (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in exacerbations, improvement in FEV1, reduction in oral corticosteroid dose). EA, CDA (reauth): Patient continues to be treated with an inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) with or without additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium) unless there is a contraindication or intolerance to these medications. CRSwNP (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in nasal polyps score [NPS, 0-8 scale], improvement in nasal congestion/obstruction score [NC, 0-3 scale]). Used in combination with another agent for CRSwNP.

## ELAPRASE (S)

---

### Products Affected

- Elaprase

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of Hunter Syndrome (Mucopolysaccharidosis II (MPS II))
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

# EMGALITY (S)

## Products Affected

- Emgality

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Episodic Migraines (EM) (120 mg/mL strength only) (initial): Diagnosis of EM. Patient has 4 to 14 migraine days per month, but no more than 14 headache days per month. Chronic Migraines (CM) (120 mg strength/mL only) (initial): Diagnosis of CM. Medication overuse headache has been considered and potentially offending medication(s) have been discontinued. Patient has greater than or equal to 15 headache days per month, of which at least 8 must be migraine days for at least 3 months. Episodic Cluster Headache (ECH) (100 mg/mL strength only) (initial): Diagnosis of episodic cluster headache. Patient has experienced at least 2 cluster periods lasting from 7 days to 365 days, separated by pain-free periods lasting at least three months. EM, CM (120 mg/mL strength only) (initial): Two of the following: a) History of failure (after at least a two month trial) or intolerance to Elavil (amitriptyline) or Effexor (venlafaxine), OR patient has a contraindication to both Elavil (amitriptyline) and Effexor (venlafaxine), b) History of failure (after at least a two month trial) or intolerance to Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate), OR patient has a contraindication to both Depakote/Depakote ER (divalproex sodium) and Topamax (topiramate), c) History of failure (after at least a two month trial) or intolerance to one of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol, OR patient has a contraindication to all of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol, or d) History of failure (after at least a two month trial) or intolerance to Atacand (candesartan), OR patient has a contraindication to Atacand (candesartan). All Indications (initial): Medication will not be used in combination with another injectable CGRP inhibitor.</p>
Age Restrictions	EM, CM, ECH (initial): 18 years of age or older.
Prescriber Restrictions	EM, CM, ECH (initial, reauth): Prescribed by or in consultation with a neurologist, headache specialist, or pain specialist.



<b>Coverage Duration</b>	EM, CM (initial): 6 months. ECH (initial): 3 months. EM, CM, ECH (reauth): 12 months.
<b>Other Criteria</b>	EM, CM (120 mg/mL strength only) (reauth): Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity. Use of acute migraine medications (e.g., non-steroidal anti-inflammatory drugs [NSAIDs] [e.g., ibuprofen, naproxen], triptans [e.g., eletriptan, rizatriptan, sumatriptan]) has decreased since the start of CGRP therapy. CM (120 mg/mL strength only) (reauth): Patient continues to be monitored for medication overuse headache. ECH (100 mg/mL strength only) (reauth): Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity. All Indications (reauthorization): Medication will not be used in combination with another injectable CGRP inhibitor.

# ENBREL (S)

## Products Affected

- Enbrel
- Enbrel Mini
- Enbrel Sureclick

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Trial and failure, contraindication, or intolerance to one disease-modifying antirheumatic drug (DMARD) [eg, methotrexate (Rheumatrex/Trexall), Arava (leflunomide), Azulfidine (sulfasalazine)]. Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Diagnosis of moderately to severely active PJIA. Trial and failure, contraindication, or intolerance to one of the following DMARDs: Arava (leflunomide) or methotrexate (Rheumatrex/Trexall). Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. Plaque psoriasis (Initial): Diagnosis of moderate to severe chronic plaque psoriasis. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. Trial and failure, contraindication, or intolerance to two NSAIDs (e.g., diclofenac, ibuprofen, meloxicam, naproxen).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RA (initial), PJIA (initial), AS (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a rheumatologist or dermatologist. Plaque Psoriasis (initial): Prescribed by or in consultation with a dermatologist.
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	RA, PJIA, PsA, AS (Reauth): Documentation of positive clinical response to therapy. Plaque psoriasis (Reauth): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline.

## ENSPRYNG (S)

---

### Products Affected

- Enspryng

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Neuromyelitis Optica Spectrum Disorder (NMOSD) (initial): Diagnosis of NMOSD. Patient is anti-aquaporin-4 (AQP4) antibody positive.
Age Restrictions	N/A
Prescriber Restrictions	NMOSD (initial): Prescribed by or in consultation with a neurologist or ophthalmologist.
Coverage Duration	NMOSD (initial, reauth): 12 months
Other Criteria	NMOSD (reauth): Documentation of positive clinical response to therapy.

## EPCLUSA PREFERRED (S)

---

### Products Affected

- Sofosbuvir/velpatasvir

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guideline. Diagnosis of chronic hepatitis C. Not used in combination with another HCV direct acting antiviral agent [e.g., Sovaldi (sofosbuvir)].
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine.
Coverage Duration	12 to 24 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline.
Other Criteria	N/A

## EPIDIOLEX (S)

---

### Products Affected

- Epidiolex

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Lennox-Gastaut syndrome (LGS): Diagnosis of seizures associated with LGS. Trial of, contraindication, or intolerance to two formulary anticonvulsants (e.g., topiramate, lamotrigine, valproate). Dravet syndrome (DS): Diagnosis of seizures associated with DS. Tuberous sclerosis complex (TSC): Diagnosis of seizures associated with TSC.
Age Restrictions	N/A
Prescriber Restrictions	LGS, DS, TSC: Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# EPOETIN ALFA (S)

## Products Affected

- Procrit

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Anemia with Chronic Kidney Disease (CKD) (Initial): Diagnosis (Dx) of CKD. Anemia by lab values (Hct less than 30% or Hgb less than 10 g/dL) collected within 30 days of request. One of the following: a) both of the following: Patient is on dialysis, patient is without ESRD OR b) all of the following: patient is not on dialysis, the rate of hemoglobin decline indicates the likelihood of requiring a red blood cell (RBC) transfusion, and reducing the risk of alloimmunization and/or other RBC transfusion-related risks is a goal. Anemia with chemo (Initial): Other causes of anemia have been ruled out. Anemia by lab values (Hct less than 30%, Hgb less than 10 g/dL) collected within the prior 2 weeks of request. Cancer is a non-myeloid malignancy. Patient is receiving chemo. Preoperative for reduction of allogeneic blood transfusion: Patient is scheduled to undergo elective, non-cardiac, non-vascular surgery. Hgb is greater than 10 to less than or equal to 13 g/dL. Patient is at high risk for perioperative transfusions. Patient is unwilling or unable to donate autologous blood pre-operatively. Anemia in hepatitis C virus (HCV)-infected pts due to ribavirin in combination with interferon/peg-interferon (Initial): Dx of HCV infection. Anemia by labs (Hct less than 36% or Hgb less than 12 g/dL) collected within 30 days of request. Patient is receiving ribavirin and one of the following: interferon alfa or peginterferon alfa. Anemia with HIV (Initial): Anemia by lab values (Hgb less than 12 g/dL or Hct less than 36%) collected within 30 days of request. Serum erythropoietin level less than or equal to 500 mU/mL. Receiving zidovudine therapy or dx of HIV. Anemia in Myelodysplastic Syndrome (MDS) (Initial): Dx of MDS. Serum erythropoietin level is 500 mU/mL or less, or dx of transfusion-dependent MDS.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A

<b>Coverage Duration</b>	CKD,HIV(Init):6mo. CKD,HIV(reauth):12mo. Chemo,HCV(all):3mo. MDS:(init) 3mo,(reauth)12mo. Preop:1mo.
<b>Other Criteria</b>	<p>Subject to ESRD review. CKD, Chemo, MDS (init): History of use or unavailability of both Aranesp and Retacrit. HIV, Preop, HCV (init): History of use or unavailability of Retacrit. CKD (Reauth): Dx of CKD. One of the following: 1) Most recent or average (avg) Hct over 3 months is 33% or less (Hgb is 11 g/dL or less) for patients on dialysis, without ESRD, 2) Most recent or avg Hct over 3 mo is 30% or less (Hgb 10 g/dL or less) for patients not on dialysis, OR 3) Most recent or avg Hct over 3 mo is 36% or less (Hgb 12 g/dL or less) for pediatric patients. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. HIV (Reauth): Most recent or avg Hct over 3 months is below 36% or most recent or avg Hgb over 3 months is below 12 g/dl. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Chemo (Reauth): Anemia by lab values (Hgb less than 10 g/dl or Hct less than 30%) collected within the prior 2 weeks of request. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Patient is receiving chemo. HCV (Reauth): Most recent or avg Hct over 3 months is 36% or less, OR most recent or avg Hgb over 3 months is 12 g/dl or less. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. If patient has demonstrated response to therapy, authorization will be issued for the full course of ribavirin therapy. MDS (Reauth): Most recent or avg Hct over 3 months is 36% or less, OR most recent or avg Hgb over 3 months is 12 g/dl or less. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Off-label uses (except MDS, HCV): Will not be approved if patient has Hgb greater than 10 g/dL or Hct greater than 30%. CKD (init, reauth), HIV (init), Chemo (init), Preop, MDS (init), HCV (init): Verify iron evaluation for adequate iron stores.</p>

## ERIVEDGE (S)

---

### Products Affected

- Erivedge

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Basal cell carcinoma: One of the following: A) Diagnosis of metastatic basal cell carcinoma OR B) Both of the following: 1) Diagnosis of locally advanced basal cell carcinoma AND 2) One of the following: a) Disease recurred following surgery or b) Patient is not a candidate for surgery and radiation.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.



## ERLEADA (S)

---

### Products Affected

- Erleada

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-metastatic castration-resistant or castration-recurrent prostate cancer (NM-CRPC): Diagnosis of non-metastatic, castration-resistant (chemical or surgical) or recurrent prostate cancer. Metastatic castration-sensitive prostate cancer (M-CSPC): Diagnosis of metastatic, castration-sensitive prostate cancer.
Age Restrictions	N/A
Prescriber Restrictions	NM-CRPC, M-CSPC: Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## ESBRIET (S)

### Products Affected

- Esbriet

- Pirfenidone

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Idiopathic pulmonary fibrosis (IPF) (initial): Diagnosis of IPF as documented by all of the following: a) exclusion of other known causes of interstitial lung disease (ILD) (eg, domestic and occupational environmental exposures, connective tissue disease, drug toxicity), AND b) one of the following: i) in patients not subjected to surgical lung biopsy, the presence of a usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) revealing IPF or probable IPF, OR ii) in patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern revealing IPF or probable IPF.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	IPF (initial): Prescribed by or in consultation with a pulmonologist
<b>Coverage Duration</b>	initial, reauth: 12 months
<b>Other Criteria</b>	IPF (reauth): Documentation of positive clinical response to therapy.

## EUCRISA (S)

---

### Products Affected

- Eucrisa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Atopic dermatitis (initial): Diagnosis of mild to moderate atopic dermatitis. Trial and failure, contraindication, or intolerance to one prescription strength topical corticosteroid, unless the affected area is sensitive (i.e., face, axillae, groin).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Reauth: Documentation of a positive clinical response to therapy (e.g., reduction in body surface area involvement, reduction in pruritus severity).

# EVRYSDI (S)

## Products Affected

- Evrysdi

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Spinal muscular atrophy (SMA) (initial): Diagnosis of spinal muscular atrophy (SMA) type I, II, or III. Both of the following: a) The mutation or deletion of genes in chromosome 5q resulting in one of the following: 1) Homozygous gene deletion or mutation (e.g., homozygous deletion of exon 7 at locus 5q13) or 2) Compound heterozygous mutation (e.g., deletion of SMN1 exon 7 [allele 1] and mutation of SMN1 [allele 2]) AND b) Patient has at least 2 copies of SMN2. Patient is not dependent on both of the following: 1) Invasive ventilation or tracheostomy and 2) Use of non-invasive ventilation beyond use for naps and nighttime sleep. At least one of the following exams (based on patient age and motor ability) has been conducted to establish baseline motor ability: Hammersmith Infant Neurological Exam (HINE) (infant to early childhood), Hammersmith Functional Motor Scale Expanded (HF MSE), Upper Limb Module (ULM) Test (Non ambulatory), or Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND), or Motor Function Measure 32 (MFM-32) Scale. Patient is not to receive concomitant chronic survival motor neuron (SMN) modifying therapy for the treatment of SMA (e.g., Spinraza). One of the following: a) patient has not previously received gene replacement therapy for the treatment of SMA (e.g., Zolgensma) or b) patient has previously received gene therapy for the treatment of SMA (e.g., Zolgensma) AND submission of medical records (e.g., chart notes) documenting that there has been an inadequate response to gene therapy (e.g., sustained decrease in at least one motor test score over a period of 6 months).</p>
<b>Age Restrictions</b>	Initial: Patient is at least 2 months of age or older
<b>Prescriber Restrictions</b>	SMA (Initial, Reauth): Prescribed by or in consultation with a neurologist with expertise in the diagnosis and treatment of SMA
<b>Coverage Duration</b>	Initial, Reauth: 12 months

<b>Other Criteria</b>	<p>SMA (Reauth): Documentation of positive clinical response to therapy. Patient (Pt) continues to not be dependent on both of the following: 1) Invasive ventilation or tracheostomy AND 2) use of non-invasive ventilation beyond use for naps and nighttime sleep. Pt is not to receive concomitant chronic survival motor neuron (SMN) modifying therapy for the treatment of SMA (e.g., Spinraza). One of the following: a) pt has not previously received gene replacement therapy for the treatment of SMA (e.g., Zolgensma) or b) pt has previously received gene therapy for the treatment of SMA (e.g., Zolgensma) AND submission of medical records (e.g., chart notes) documenting that there has been an inadequate response to gene therapy (e.g., sustained decrease in at least one motor test score over a period of 6 months).</p>
-----------------------	--

## EXKIVITY (S)

### Products Affected

- Exkivity

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of non-small cell lung cancer (NSCLC). Disease is one of the following: a) locally advanced or b) metastatic. Disease is epidermal growth factor receptor (EGFR) exon 20 insertion mutation positive as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient has progressed on or following prior treatment with a platinum-containing regimen (e.g., carboplatin, cisplatin).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## FARYDAK (S)

---

### Products Affected

- Farydak

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple Myeloma (MM): Diagnosis of MM. Used in combination with both of the following: Velcade (bortezomib) and dexamethasone. Patient has received at least two prior treatment regimens which included both of the following: Velcade (bortezomib) and an immunomodulatory agent [eg, Revlimid (lenalidomide), Thalomid (thalidomide)].
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist/hematologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# FASENRA (S)

## Products Affected

- Fasenra

- Fasenra Pen

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Asthma (initial): Diagnosis of severe asthma. Asthma is an eosinophilic phenotype as defined by a baseline (pre-treatment) peripheral blood eosinophil level greater than or equal to 150 cells per microliter. One of the following: 1) Patient has had at least one or more asthma exacerbations requiring systemic corticosteroids (e.g., prednisone) within the past 12 months, OR 2) Any prior intubation for an asthma exacerbation, OR 3) Prior asthma-related hospitalization within the past 12 months. Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) High-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist (e.g., montelukast), long-acting beta-2 agonist (LABA) (e.g., salmeterol), tiotropium], OR b) One maximally-dosed combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone/vilanterol)].
Age Restrictions	Asthma (Initial): Patient is 12 years of age or older
Prescriber Restrictions	Asthma (Initial/Reauth): Prescribed by or in consultation with a pulmonologist or allergist/immunologist
Coverage Duration	Asthma (init): 6 months. Asthma (reauth): 12 months



<b>Other Criteria</b>	<p>Asthma (Reauth): Documentation of positive clinical response to therapy (e.g., reduction in exacerbations, improvement in forced expiratory volume in 1 second [FEV1], decreased use of rescue medications). Patient continues to be treated with an inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) with or without additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium) unless there is a contraindication or intolerance to these medications.</p>
-----------------------	--

## FENTANYL (S)

### Products Affected

- Fentanyl Citrate Oral Transmucosal

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the management of breakthrough cancer pain. Patient is currently taking a long-acting opioid around the clock for cancer pain. Patient must have at least a one week history of ONE of the following medications to demonstrate tolerance to opioids: Morphine sulfate at doses of greater than or equal to 60 mg/day, Fentanyl transdermal patch at doses greater than or equal to 25 µg/hr, Oxycodone at a dose of greater than or equal to 30 mg/day, Oral hydromorphone at a dose of greater than or equal to 8 mg/day, Oral oxymorphone at a dose of greater than or equal to 25 mg/day, or an alternative opioid at an equianalgesic dose (e.g., oral methadone greater than or equal to 20 mg/day).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with one of the following: Pain specialist, Oncologist, Hematologist, Hospice care specialist, or Palliative care specialist.
Coverage Duration	12 months
Other Criteria	N/A

# FERRIPROX (S)

## Products Affected

- Deferiprone

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Transfusional iron overload (Initial): Diagnosis of transfusional iron overload due to one of the following: thalassemia syndromes, sickle cell disease, or other transfusion-dependent anemias. Patient has Absolute Neutrophil Count (ANC) greater than $1.5 \times 10^9/L$ . One of the following: A) Trial and failure, defined as serum ferritin greater than 2,500 mcg/L, to one chelation therapy (i.e., deferoxamine, deferasirox) OR B) History of contraindication or intolerance to one chelation therapy (i.e., deferoxamine, deferasirox).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	All uses (reauth): Documentation of positive clinical response to therapy (e.g., greater than or equal to 20% decline in serum ferritin levels from baseline, decrease in liver iron concentration). ANC greater than $1.5 \times 10^9/L$ .

## FINTEPLA (S)

---

### Products Affected

- Fintepla

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of seizures associated with Dravet syndrome
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## FIRAZYR (S)

### Products Affected

- Icatibant Acetate
- Sajazir

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Treatment of hereditary angioedema (HAE) attacks: Diagnosis of HAE. Diagnosis has been confirmed by C1 inhibitor (C1-INh) deficiency or dysfunction (Type I or II HAE) as documented by one of the following: a) C1-INH antigenic level below the lower limit of normal OR b) C1-INH functional level below the lower limit of normal. For the treatment of acute HAE attacks. Not used in combination with other approved treatments for acute HAE attacks.
Age Restrictions	N/A
Prescriber Restrictions	HAE: Prescribed by or in consultation with an immunologist or an allergist
Coverage Duration	12 months
Other Criteria	N/A

## FIRMAGON (S)

---

### Products Affected

- Firmagon INJ 120MG/VIAL, 80MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of advanced or metastatic prostate cancer.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## FOTIVDA (S)

---

### Products Affected

- Fotivda

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of renal cell carcinoma. Disease is one of the following: relapsed or refractory. Patient has received two or more prior systemic therapies (e.g., cabozantinib + nivolumab, lenvatinib + pembrolizumab, etc.).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, nephrologist, or urologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## GALAFOLD (S)

---

### Products Affected

- Galafold

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Fabry Disease (FD) (initial): Diagnosis of FD. Patient has an amenable galactosidase alpha gene (GLA) variant based on in vitro assay data. FD (initial, reauthorization): Will not be used in combination with Fabrazyme (agalsidase beta).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	FD (initial, reauth): 12 months.
Other Criteria	FD (reauthorization): Documentation of positive clinical response to therapy.



## GAMASTAN S/D (S)

### Products Affected

- Gamastan

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication).
Required Medical Information	Immune globulin is being used intramuscularly. The immune globulin will be administered at the minimum effective dose and appropriate frequency for the prescribed diagnosis. Patient requires immunization for hepatitis A, measles, rubella, or varicella.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months (Approve one dose only)
Other Criteria	Subject to Part B vs D review.

# GAVRETO (S)

## Products Affected

- Gavreto

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Presence of metastatic rearranged during transfection (RET) gene fusion-positive tumor(s). Medullary Thyroid Cancer (MTC): Diagnosis of medullary thyroid cancer (MTC). Disease is one of the following: advanced or metastatic. Disease has presence of rearranged during transfection (RET) gene mutation tumor(s). Disease requires treatment with systemic therapy. Thyroid Cancer: Diagnosis of thyroid cancer. Disease is one of the following: advanced or metastatic. Disease has presence of rearranged during transfection (RET) gene fusion-positive tumor(s). Disease requires treatment with systemic therapy. One of the following: patient is radioactive iodine-refractory or radioactive iodine therapy is not appropriate.
Age Restrictions	N/A
Prescriber Restrictions	NSCLC, MTC: Prescribed by or in consultation with an oncologist. Thyroid Cancer: Prescribed by or in consultation with an endocrinologist or an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy

## GILENYA (S)

### Products Affected

- Gilenya

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS) (initial): Diagnosis of a relapsing form of MS (eg, clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	MS (initial, reauth): Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	MS (initial, reauth): 12 months
<b>Other Criteria</b>	MS (reauth): Documentation of positive clinical response to therapy (e.g., stability in radiologic disease activity, clinical relapses, disease progression).

# GILOTRIF (S)

## Products Affected

- Gilotrif

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): A) Diagnosis of advanced or metastatic (stage IIIB or IV) NSCLC AND B) One of the following: 1) Both of the following: a) Tumors have non-resistant epidermal growth factor (EGFR) mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) AND b) GILOTRIF will be used as first-line treatment, OR 2) All of the following: a) disease progressed after platinum-based chemotherapy and b) squamous NSCLC.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# GLATIRAMER ACETATE (S)

## Products Affected

- Glatiramer Acetate

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS) (initial): Diagnosis of a relapsing form of MS (eg, clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	MS (initial, reauth): Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	MS (initial, reauth): 12 months
<b>Other Criteria</b>	MS (reauth): Documentation of positive clinical response to therapy (e.g., stability in radiologic disease activity, clinical relapses, disease progression).

# GLEEVEC (S)

## Products Affected

- Imatinib Mesylate

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	One of the following: A) Diagnosis of Philadelphia chromosome positive (Ph+)/BCR ABL-positive chronic myelogenous leukemia (CML) OR B) Ph+/BCR ABL+ acute lymphoblastic leukemia (ALL) OR C) Gastrointestinal stromal tumor (GIST) AND one of the following: 1) Patient has documented c-KIT (CD117) positive unresectable or metastatic malignant GIST, OR 2) Patient had resection of c-KIT (CD117) positive GIST and imatinib will be used as adjuvant therapy OR D) Dermatofibrosarcoma protuberans that is unresectable, recurrent, or metastatic OR E) Hypereosinophilic syndrome or chronic eosinophilic leukemia OR F) Myelodysplastic syndrome (MDS) or myeloproliferative disease associated with platelet-derived growth factor receptor gene rearrangements OR G) Aggressive systemic mastocytosis without the D816V c-KIT mutation or with c-KIT mutational status unknown.
Age Restrictions	N/A
Prescriber Restrictions	Hypereosinophilic syndrome or chronic eosinophilic leukemia, Aggressive systemic mastocytosis: Prescribed by or in consultation with an oncologist, hematologist, allergist, or immunologist. All other uses: Prescribed by or in consultation with an oncologist or hematologist.
Coverage Duration	All uses: 12 months
Other Criteria	All uses: Approve for continuation of prior therapy.

## GROWTH HORMONE, PREFERRED (S)

### Products Affected

- Genotropin

- Genotropin Miniquick

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>PGHD(initial):less than 4mo w/suspected GD based on clinical presentation (eg, persistent neonatal hypoglycemia, persistent/prolonged neonatal jaundice/elev bilirubin, male infant with microgenitalia, midline anatomical defects, failure to thrive),OR hx neonatal hypoglycemia assoc w/pituitary dz,or panhypopituitarism dx,or all of the following: PGHD dx [confrmd by ht (utilizing age and gender grwth charts related to ht) documented(doc) by ht more than 2.0SD below midparental ht or more than 2.25SD below population(pop) mean (below 1.2 percentile for age and gender),or grwth velocity more than 2SD below mean for age and gender, or delayed skeletal maturation more than 2SD below mean for age and gender (eg,delayed more than 2yrs compared w/chronological age)].</p> <p>PWS(reauth):evidence of positive response to tx(eg,incr in total LBM, decr in fat mass) and expctd adult ht not attained and doc of expctd adult ht goal.</p> <p>GFSGA(initial):SGA dx based on catchup grwth failure in 1st 24mo of life using 0-36mo grwth chart confrmd by birth wt or length below 3rd percentile for gestational age(more than 2SD below pop mean) and ht remains at or below 3rd percentile (more than 2SD below pop mean).</p> <p>TS,NS(initial):ped grwth failure dx assoc w/TS w/doc female w/bone age less than 14yrs, or NS and ht below 5th percentile on grwth charts for age and gender.</p> <p>SHOX(initial):ped grwth failure dx w/SHOX gene deficiency confirmed by genetic testing.</p> <p>GFCRI(initial): ped grwth failure dx assoc w/CRI.</p> <p>ISS(initial):ISS dx, diagnostic eval excluded other causes assoc w/short stature(eg GHD, chronic renal insufficiency), doc ht at or below -2.25SD score below corresponding mean ht for age and gender assoc with growth rates unlikely to permit attainment of adult height in the normal range.</p> <p>PGHD,NS,SHOX,GFCRI,ISS (initial): doc male w/bone age less than 16yrs or female w/bone age less than 14yrs.</p> <p>PGHD,GFSGA,TS/NS,SHOX,GFCRI,ISS(reauth):expctd adult ht not attained and doc of expctd adult ht goal.</p>

<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PGHD, PWS, GFSGA, TS/NS, SHOX, AGHD, TPAP, IGHDA, ISS: prescribed by endocrinologist. GFCRI: prescribed by endocrinologist or nephrologist
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months



Other Criteria	<p>AGHD(initial):dx of AGHD with clin records supporting dx of childhood-onset GHD, or adult-onset GHD w/clin records doc hormone deficiency d/t hypothalamic-pituitary dz from organic or known causes (eg,damage from surgery, cranial irradiation, head trauma, subarachnoid hemorrhage) and pt has 1GH stim test (insulin tolerance test [ITT], arginine/GHRH,glucagon,arginine,macimorelin) to confirm adult GHD w/peak GH values ([ITT at or below 5mcg/L],[GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2],[glucagon at or below 3mcg/L],[Arg at or below 0.4mcg/L],[macimorelin less than 2.8 ng/mL 30, 45, 60 and 90 mins after admin]) or doc deficiency of 3 anterior pituitary hormones (prolactin,ACTH,TSH,FSH/LH) and IGF-1/somatomedinC below age and gender adjstd nrml range as provided by physicians lab.</p> <p>AGHD,IGHDA(reauth):monitoring as demonstrated by doc w/in past 12mo of IGF-1/somatomedinC level. TransitionPhaseAdolescent</p> <p>Pts(TPAP)(initial): attained expctd adult ht or closed epiphyses on bone radiograph, and doc high risk of GHD d/t GHD in childhood (from embryopathic/congenital defects, genetic mutations, irreversible structural hypothalamic-pituitary dz, panhypopituitarism, or deficiency of 3 anterior pituitary hormones: ACTH,TSH,prolactin,FSH/LH), w/IGF-1/somatomedinC below age and gender adj nrml range as provided by physicians lab, or pt does not have low IGF-1/somatomedinC and d/c GH tx for at least 1mo, and pt has 1 GH stim test (ITT,GHRH+ARG,ARG,glucagon,macimorelin) after d/c of tx for at least 1mo w/peak GH value [ITT at or below 5mcg/L], [GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2], [glucagon at or below 3mcg/L], [Arg at or below 0.4mcg/L], [macimorelin less than 2.8 ng/mL 30, 45, 60 and 90 mins after admin], or at low risk of severe GHD(eg d/t isolated and/or idiopathic GHD) and d/c GH tx for at least 1mo, and pt has 1 GH stim test (ITT, GHRH+ARG, ARG, glucagon, macimorelin) after d/c of tx for at least 1mo w/corresponding peak GH value [ITT at or below 5mcg/L], [GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2], [glucagon at or below 3mcg/L], [Arg at or below 0.4mcg/L], [macimorelin less than 2.8 ng/mL 30, 45, 60 and 90 mins after admin]. TPAP(reauth): evidence of positive response to therapy (eg,incr in total lean body mass, exercise capacity or IGF-1 and IGFBP-3). IGHDA(initial):doc GHD after 2 GH stim tests(ITT,L-ARG,glucagon,macimorelin), w/ 2 corresponding peak GH values [ITT at or below 5mcg/L],[Arg at or below 5mcg/L],[glucagon at or below 3mcg/L],[macimorelin less than 2.8 ng/mL 30,45,60,90 mins after admin].</p>
----------------	---

# HERZUMA (S)

## Products Affected

- Herzuma INJ 420MG

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast cancer: Diagnosis of human epidermal growth factor receptor 2 (HER2)-overexpressing breast cancer. One of the following treatment regimens: a) As adjuvant treatment, b) metastatic disease and one of the following: 1) used in combination with a taxane (eg, docetaxel, paclitaxel), or 2) used as a single agent in a patient who has received one or more chemotherapy regimens for metastatic disease, or c) used in combination with Perjeta (pertuzumab). Gastric Cancer: Diagnosis of HER2-overexpressing gastric or gastroesophageal junction adenocarcinoma (locally advanced, recurrent, or metastatic). Used in combination with one of the following treatment regimens: a) Platinol (cisplatin) and Adrucil (5-fluorouracil), or b) Platinol (cisplatin) and Xeloda (capecitabine). All indications: One of the following: trial and failure, or intolerance to both Kanjinti (trastuzumab-anns) and Trazimera (trastuzumab-qyyp), OR approve for continuation of prior therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## HRM - ANTIPSYCHOTICS

---

### Products Affected

- Thioridazine Hcl TABS 100MG, 10MG, 25MG, 50MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The drug is being used for an FDA-approved diagnosis AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication. Trial and failure, contraindication or intolerance to one of the following: haloperidol, fluphenazine, or an atypical antipsychotic.
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Applies to New Starts only.

## HRM - MEGESTROL SUSPENSION

---

### Products Affected

- Megestrol Acetate SUSP 40MG/ML, 625MG/5ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

# HRM - MEGESTROL TABLET

---

## Products Affected

- Megestrol Acetate TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Applies to New Starts only.

## HRM - PHENOBARBITAL, PENTOBARBITAL

---

### Products Affected

- Phenobarbital ELIX 20MG/5ML
- Phenobarbital TABS 100MG, 15MG, 16.2MG, 30MG, 32.4MG, 60MG, 64.8MG, 97.2MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The drug is being used for an FDA-approved diagnosis AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Applies to New Starts only.

## HRM - SKELETAL MUSCLE RELAXANTS

---

### Products Affected

- Chlorzoxazone TABS 500MG
- Cyclobenzaprine Hydrochloride TABS 10MG, 5MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication.
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

## HRM - TCA

---

### Products Affected

- Amitriptyline Hcl TABS 100MG, 150MG, 75MG
- Amitriptyline Hydrochloride TABS 10MG, 25MG, 50MG
- Doxepin Hcl CAPS 75MG
- Doxepin Hcl CONC
- Doxepin Hydrochloride CAPS 100MG, 10MG, 150MG, 25MG, 50MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Applies to New Starts only.



# HUMIRA (S)

## Products Affected

- Humira
- Humira Pediatric Crohns Disease Starter Pack
- Humira Pen
- Humira Pen-cd/uc/hs Starter
- Humira Pen-pediatric Uc Starter Pack
- Humira Pen-ps/uv Starter

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Trial and failure, contraindication, or intolerance (TF/C/I) to one disease modifying antirheumatic drug (DMARD) [eg, methotrexate (Rheumatrex/Trexall), Arava (leflunomide), Azulfidine (sulfasalazine)].</p> <p>Juvenile Idiopathic Arthritis (JIA) (Initial): Diagnosis of moderately to severely active polyarticular JIA. TF/C/I to one of the following DMARDs: Arava (leflunomide) or methotrexate (Rheumatrex/Trexall).</p> <p>Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. Plaque psoriasis (Initial): Diagnosis of moderate to severe chronic plaque psoriasis. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. TF/C/I to two NSAIDs (e.g., diclofenac, ibuprofen, meloxicam, naproxen).</p> <p>Crohn's Disease (CD) (Initial): Diagnosis of moderately to severely active CD. TF/C/I to one of the following conventional therapies: 6-mercaptopurine, Imuran (azathioprine), corticosteroid (eg, prednisone, methylprednisolone), methotrexate (Rheumatrex/Trexall).</p> <p>Ulcerative Colitis (UC) (Initial): Diagnosis of moderately to severely active UC. TF/C/I to one of the following conventional therapies: 6-mercaptopurine, Imuran (azathioprine), corticosteroid (eg, prednisone, methylprednisolone), aminosalicylate [eg, mesalamine (Asacol, Pentasa, Rowasa), Dipentum (olsalazine), Azulfidine (sulfasalazine)].</p> <p>Hidradenitis suppurativa (Initial): Diagnosis of moderate to severe hidradenitis suppurativa (ie, Hurley Stage II or III). Uveitis (initial): Diagnosis of non-infectious uveitis, classified as intermediate, posterior, or panuveitis.</p>
Age Restrictions	N/A

<b>Prescriber Restrictions</b>	RA, AS, JIA (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. Plaque Psoriasis, HS (initial): Prescribed by or in consultation with a dermatologist. CD, UC (initial): Prescribed by or in consultation with a gastroenterologist. Uveitis (initial): Prescribed by or in consultation with an ophthalmologist or rheumatologist.
<b>Coverage Duration</b>	UC (Initial): 12 wks. UC (reauth): 12 mo. All other indications (initial, reauth): 12 mo.
<b>Other Criteria</b>	RA, JIA, PsA, AS, CD, Hidradenitis suppurativa (HS), Uveitis (Reauth): Documentation of positive clinical response to therapy. Plaque psoriasis (Reauth): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline. UC (Reauth): For patients who initiated therapy within the past 12 weeks: Documentation of clinical remission or significant clinical benefit by eight weeks (Day 57) of therapy OR For patients who have been maintained on therapy for longer than 12 weeks: Documentation of positive clinical response to therapy.

## IBRANCE (S)

### Products Affected

- Ibrance

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast Cancer: Diagnosis of advanced or metastatic breast cancer. Disease is a) hormone receptor (HR)-positive, and b) human epidermal growth factor receptor 2 (HER2)-negative. One of the following: 1) used in combination with an aromatase inhibitor (e.g., anastrozole, letrozole, exemestane) and one of the following: a) patient is a male, or b) patient is a postmenopausal woman, OR 2) both of the following: used in combination with Faslodex (fulvestrant) and disease has progressed following endocrine therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

# ICLUSIG (S)

## Products Affected

- Iclusig

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic myelogenous leukemia: Diagnosis of chronic myelogenous leukemia AND One of the following: A) Trial and failure, contraindication, or intolerance to at least TWO other tyrosine kinase inhibitors (e.g., Gleevec [imatinib], Sprycel, Tasigna, and Bosulif) or B) Patient has the T315I mutation. Acute Lymphoblastic Leukemia: Diagnosis of Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) AND One of the following: A) Trial and failure, resistance relapse, contraindication, or intolerance to at least TWO other FDA-approved tyrosine kinase inhibitors (e.g., Gleevec [imatinib], Sprycel), or B) Patient has the T315I mutation.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist or hematologist
<b>Coverage Duration</b>	All uses: 12 months
<b>Other Criteria</b>	All uses: Approve for continuation of prior therapy.

## IDHIFA (S)

### Products Affected

- Idhifa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute Myeloid Leukemia (AML): Diagnosis of AML. Disease is relapsed or refractory. Patient has an isocitrate dehydrogenase-2 (IDH2) mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., Abbott RealTime IDH2 assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist/oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## ILUMYA (S)

### Products Affected

- Ilumya

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Plaque Psoriasis (initial): Diagnosis of moderate-to-severe plaque psoriasis. Trial and failure, contraindication, or intolerance to two of the following: Enbrel (etanercept), Humira (adalimumab), Skyrizi (risankizumab), Cosentyx (secukinumab), or for continuation of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	Plaque Psoriasis (initial): Prescribed by or in consultation with a dermatologist
Coverage Duration	Initial, reauth: 12 months
Other Criteria	Plaque Psoriasis (Reauth): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline.

# IMBRUVICA (S)

## Products Affected

- Imbruvica

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Mantle cell lymphoma (MCL): Diagnosis of MCL AND patient has relapsed or is refractory to at least one prior therapy for the treatment of MCL. Chronic lymphocytic leukemia (CLL): Diagnosis of CLL. Waldenstrom's macroglobulinemia: Diagnosis of Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma. Small lymphocytic lymphoma (SLL): Diagnosis of SLL. Marginal zone lymphoma (MZL): Diagnosis of MZL AND patient has received at least one prior anti-CD20-based therapy for MZL [e.g., Rituxan (rituximab), Zevalin (ibritumomab), Gazyva (obinutuzumab, etc.)]. Chronic graft versus host disease (cGVHD): Diagnosis of cGVHD AND trial and failure of one or more lines of systemic therapy (e.g., corticosteroids like prednisone or methylprednisolone, mycophenolate).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses (except chronic graft versus host disease): Prescribed by or in consultation with an oncologist or hematologist. Chronic graft versus host disease: Prescribed by or in consultation with a hematologist, oncologist, or physician experienced in the management of transplant patients.
<b>Coverage Duration</b>	All Uses: 12 months
<b>Other Criteria</b>	All Uses: Approve for continuation of prior therapy.

## INCRELEX (S)

### Products Affected

- Increlex

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Insulin-like Growth Factor-1 (IGF-1) deficiency (initial): Diagnosis of severe primary IGF-1 deficiency. Height standard deviation score of -3.0 or less. Basal IGF-1 standard deviation score of -3.0 or less. Normal or elevated growth hormone (GH). GH gene deletion (initial): Diagnosis of GH gene deletion in patients who have developed neutralizing antibodies to GH.
Age Restrictions	N/A
Prescriber Restrictions	Initial: Prescribed by or in consultation with an endocrinologist
Coverage Duration	Initial, reauth: 12 months
Other Criteria	(Reauth): Documentation of positive clinical response to therapy.



## INLYTA (S)

### Products Affected

- Inlyta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Renal cell carcinoma (RCC): Diagnosis of RCC. One of the following: (1) disease has relapsed or (2) diagnosis of stage IV disease.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## INQOVI (S)

### Products Affected

- Inqovi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Myelodysplastic syndrome (MDS): Diagnosis of myelodysplastic syndrome. Patient has ONE of the following French- American-British subtypes: a) refractory anemia, b) refractory anemia with ringed sideroblasts, c) refractory anemia with excess blasts, or d) chronic myelomonocytic leukemia (CMML).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## INREBIC (S)

### Products Affected

- Inrebic

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Myelofibrosis: Diagnosis of one of the following: primary myelofibrosis, post-polycythemia vera myelofibrosis, or post-essential thrombocythemia myelofibrosis.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# INTRON A (S)

## Products Affected

- Intron A

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic hepatitis B: Diagnosis of chronic hepatitis B infection, and patient is without decompensated liver disease. Chronic Hepatitis C: Diagnosis of chronic hepatitis C, patient without decompensated liver disease, patient has not previously been treated with interferon, and one of the following - 1) used in combination with ribavirin or 2) contraindication or intolerance to ribavirin. Metastatic renal cell carcinoma (RCC): diagnosis of metastatic RCC, used in combination with Avastin (bevacizumab). Other: diagnosis of condylomata acuminata (genital or perianal), diagnosis of hairy cell leukemia, diagnosis of AIDS-related Kaposi's sarcoma, diagnosis of malignant melanoma, diagnosis of Stage III or IV follicular Non-Hodgkin's Lymphoma.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RCC: Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	HepB, HepC: 48 wks. Condylomata acuminata (genital or perianal): 6 wks. Other: 12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## IRESSA (S)

### Products Affected

- Iressa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC AND Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## ISOTRETINOIN (S)

### Products Affected

- Amnesteem
- Claravis
- Isotretinoin CAPS 10MG, 20MG, 30MG, 40MG
- Myorisan
- Zenatane

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Acne (initial): Diagnosis of acne. One of the following: A) Prescribed by a dermatologist or, B) Trial and failure, contraindication or intolerance to an adequate trial (at least 6 weeks) on two of the following conventional therapy regimens: a) topical retinoid or retinoid-like agent [eg, Retin-A/Retin-A Micro (tretinoin)], b) oral antibiotic [eg, Ery-Tab (erythromycin), Minocin (minocycline)], c) topical antibiotic with or without benzoyl peroxide [eg, Cleocin-T (clindamycin), erythromycin, BenzaClin (benzoyl peroxide/clindamycin), Benzamycin (benzoyl peroxide/erythromycin)].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Acne (initial): 5 months. Acne (reauth): Retreatment - 5 months, Dose Titration - 1 month
<b>Other Criteria</b>	Acne, Retreatment (reauth): After more than 2 months off therapy, persistent or recurring acne is still present. Acne, Dose Titration (reauth): Confirmation that the total cumulative dose is less than 150 mg/kg.

## ISTODAX (S)

### Products Affected

- Romidepsin INJ 27.5MG/5.5ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cutaneous T-cell lymphoma (CTCL): Diagnosis of CTCL. Trial and failure, contraindication, or intolerance to at least one systemic therapy for the treatment of CTCL [e.g., Trexall (methotrexate), Targretin (bexarotene), cyclophosphamide] Peripheral T-cell lymphoma (PTCL): Diagnosis of PTCL. Trial and failure, contraindication, or intolerance to at least one therapy for the treatment of PTCL (e.g., conventional chemotherapy such as CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone), CHOEP (cyclophosphamide, doxorubicin, vincristine, etoposide, prednisone), Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin), HyperCVAD (cyclophosphamide, vincristine, doxorubicin, dexamethasone) alternating with high-dose methotrexate and cytarabine).
Age Restrictions	N/A
Prescriber Restrictions	CTCL, PTCL: Prescribed by or in consultation with an oncologist/hematologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## ISTURISA (S)

### Products Affected

- Isturisa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cushing's disease (initial): Diagnosis of Cushing's disease. One of the following: a) Patient is not a candidate for pituitary surgery, OR b) Pituitary surgery has not been curative for the patient.
Age Restrictions	N/A
Prescriber Restrictions	Cushing's disease (initial): Prescribed by or in consultation with an endocrinologist.
Coverage Duration	Cushing's disease (initial, reauth): 12 months
Other Criteria	Cushing's disease (reauth): Documentation of positive clinical response to therapy (e.g., a clinically meaningful reduction in 24-hour urinary free cortisol levels, improvement in signs or symptoms of the disease).



## IVERMECTIN (S)

### Products Affected

- Ivermectin TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Strongyloidiasis: Diagnosis of intestinal (i.e., nondisseminated) strongyloidiasis due to the nematode parasite Strongyloides stercoralis OR both of the following: member received the drug within the past 120 days and member requires continuation of therapy. Onchocerciasis: Diagnosis of onchocerciasis due to the nematode parasite Onchocerca volvulus OR both of the following: member received the drug within the past 120 days and member requires continuation of therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Strongyloidiasis: 3 weeks. Onchocerciasis: 6 months.
Other Criteria	N/A

## IVIG (S)

### Products Affected

- Bivigam INJ 5GM/50ML
- Gammagard S/d Iga Less Than 1mcg/ml INJ 5GM

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	All uses (initial, reauth): Contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication).
Required Medical Information	Initial: Immune globulin (Ig) will be administered at the minimum effective dose and appropriate frequency for the prescribed diagnosis. For IVIG – Ig is being used intravenously (IV) AND One of the following diagnoses: [A] Primary Immunodeficiency 1) Common variable immunodeficiency. 2) Congenital agammaglobulinemia (X-linked or autosomal recessive). 3) Severe combined immunodeficiencies. 4) Wiskott-Aldrich syndrome. OR 5) Other PI with an immunologic evaluation including IgG levels below the normal laboratory value for the patient's age at the time of diagnosis and the patient lacks an adequate response to protein and polysaccharide antigens (i.e., tetanus toxoid or diphtheria toxoid and pneumovax or HiB vaccine). [B] Secondary Acquired Antibody Deficiency 1) B-cell chronic lymphocytic leukemia with an Ig level less than 500 mg/dL OR history of recurrent bacterial infections. 2) HIV infection with an Ig level less than 400 mg/dL OR Patient has active bleeding or a platelet count less than $10 \times 10^9/L$ . 3) Multiple myeloma in plateau phase and patient has hypogammaglobulinemia. [C] Hematological Autoimmune Disorders 1) Acquired (pure) red cell aplasia (PRCA) that is immunologic and patient had a trial and failure, contraindication, or intolerance (TF/C/I) to a corticosteroid and an immunosuppressant (i.e., cyclophosphamide, cyclosporine) OR patient has viral PRCA caused by parvovirus B19. 2) Fetal alloimmune thrombocytopenia. 3) Hemolytic disease of the newborn and the patient has established hyperbilirubinemia. 4) Idiopathic thrombocytopenic purpura and patient had a TF/C/I to a corticosteroid OR a platelet count less than 30,000 cells/mm <sup>3</sup> . Continued in Other Criteria Section.
Age Restrictions	HIV (initial): patient is less than or equal to 12 years of age.

<b>Prescriber Restrictions</b>	All uses (initial, reauth): Prescribed by or in consultation with a physician who has specialized expertise in managing patients on immune globulin therapy (e.g., immunologist, hematologist, neurologist).
<b>Coverage Duration</b>	4 months: Solid organ transplant. 12 months: all other diagnoses.
<b>Other Criteria</b>	<p>[D] Neuromuscular Autoimmune Disorders 1) Chronic inflammatory demyelinating polyneuropathy. 2) Guillain-Barré syndrome. 3) Inflammatory myopathies (dermatomyositis or polymyositis) AND Patient had a TF/C/I to a corticosteroid AND an immunosuppressant (i.e., azathioprine, methotrexate, cyclosporine A, cyclophosphamide, or tacrolimus). 4) Lambert-Eaton myasthenic syndrome AND Patient had a TF/C/I to a corticosteroid AND an immunosuppressant (e.g., azathioprine). 5) Multifocal motor neuropathy. 6) Myasthenia gravis with severe exacerbations or myasthenic crises AND Patient had a TF/C/I to a corticosteroid AND an immunosuppressant (i.e., azathioprine, cyclosporine, cyclophosphamide, or mycophenolate mofetil). 7) Stiff person syndrome AND Patient had a TF/C/I to at least 2 standard therapies (i.e., benzodiazepines, muscle relaxants, or anti-convulsants).</p> <p>[E] Other Disorders 1) Autoimmune blistering disease AND Patient had a TF/C/I to a corticosteroid AND an immunosuppressant (i.e., cyclophosphamide, dapsone, methotrexate, azathioprine, or mycophenolate mofetil). 2) Kawasaki syndrome. 3) Solid organ transplant and IVIG is being used for CMV prophylaxis, or patient is a kidney transplant recipient and has donor specific antibodies, or patient has steroid-resistant rejection and had a TF/C/I to standard therapies. For SCIG (Gamunex-C, Gammagard Liquid, Gammaked only)- Immune globulin is being used subcutaneously AND One of the following PI diagnoses: 1) Common variable immunodeficiency. 2) Congenital agammaglobulinemia (X-linked or autosomal recessive). 3) Severe combined immunodeficiencies. 4) Wiskott-Aldrich syndrome. OR 5) Other PI with an immunologic evaluation including IgG levels below the normal laboratory value for the patient's age at the time of diagnosis and patient lacks an adequate response to protein and polysaccharide antigens (i.e., tetanus toxoid or diphtheria toxoid and pneumovax or HiB vaccine). All products: Subject to Part B vs. Part D review. For non-oncology renewal, the patient has experienced an objective improvement on immune globulin therapy and the immune globulin will be administered at the minimum effective dose (by decreasing the dose, increasing the frequency, or implementing both strategies) for maintenance therapy.</p>

## JAKAFI (S)

### Products Affected

- Jakafi

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Myelofibrosis: Diagnosis of primary myelofibrosis, OR post-polycythemia vera myelofibrosis, OR post-essential thrombocythemia myelofibrosis. Polycythemia vera: Diagnosis of polycythemia vera, AND trial and failure, contraindication, or intolerance to hydroxyurea. Acute graft versus host disease (aGVHD): Diagnosis of aGVHD. Disease is steroid-refractory. Chronic graft versus host disease (cGVHD): Diagnosis of cGVHD. Trial and failure of at least one or more lines of systemic therapy (e.g., corticosteroids, mycophenolate, etc.).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Myelofibrosis, Polycythemia vera: Prescribed by or in consultation with a hematologist/oncologist. aGVHD, cGVHD: Prescribed by or in consultation with one of the following: hematologist, oncologist, physician experienced in the management of transplant patients.
<b>Coverage Duration</b>	12 months.
<b>Other Criteria</b>	Approve for continuation of prior therapy.

# JUXTAPID (S)

## Products Affected

- Juxtapid

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Homozygous familial hypercholesterolemia (HoFH) (initial): Submission of medical records (eg, chart notes, laboratory values) documenting diagnosis of HoFH as confirmed by one of the following: a) genetic confirmation of 2 mutations in the LDL receptor, ApoB, PCSK9, or LDL receptor adaptor protein 1 (ie, LDLRAP1 or ARH), or b) both of the following: 1) either untreated LDL-C greater than 500 mg/dL or treated LDL-C greater than 300 mg/dL AND 2) either xanthoma before 10 years of age or evidence of heterozygous FH in both parents. Patient is receiving other lipid-lowering therapy. Trial and failure, contraindication, or intolerance to Repatha therapy. Not used in combination with a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	HoFH (initial, reauth): Prescribed by a cardiologist, endocrinologist, or lipid specialist.
<b>Coverage Duration</b>	HoFH (initial): 6 months. (reauth): 12 months
<b>Other Criteria</b>	HoFH (reauthorization): Patient continues to receive other lipid-lowering therapy. Submission of medical records (eg, chart notes, laboratory values) documenting LDL-C reduction while on therapy. Not used in combination with a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor.

# KALYDECO (S)

## Products Affected

- Kalydeco

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystic Fibrosis (CF) (Initial): Diagnosis of cystic fibrosis. Patient has at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	CF (Initial): 4 months of age or older
<b>Prescriber Restrictions</b>	CF (initial): Prescribed by or in consultation with a specialist affiliated with a CF care center or pulmonologist
<b>Coverage Duration</b>	CF (initial, reauth): 12 months
<b>Other Criteria</b>	CF (Reauth): Documentation of positive clinical response (i.e. improvement in lung function [percent predicted forced expiratory volume in one second (PPFEV1)], decreased number of pulmonary exacerbations) while on therapy.

# KANJINTI (S)

## Products Affected

- Kanjinti

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast cancer: Diagnosis of human epidermal growth factor receptor 2 (HER2)-overexpressing breast cancer. One of the following treatment regimens: a) As adjuvant treatment, b) metastatic disease and one of the following: 1) used in combination with a taxane (eg, docetaxel, paclitaxel), or 2) used as a single agent in a patient who has received one or more chemotherapy regimens for metastatic disease, or c) used in combination with Perjeta (pertuzumab). Gastric Cancer: Diagnosis of HER2-overexpressing gastric or gastroesophageal junction adenocarcinoma (locally advanced, recurrent, or metastatic). Used in combination with one of the following treatment regimens: a) Platinol (cisplatin) and Adrucil (5-fluorouracil), or b) Platinol (cisplatin) and Xeloda (capecitabine).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## KANUMA (S)

---

### Products Affected

- Kanuma

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Lysosomal acid lipase deficiency: Diagnosis of lysosomal acid lipase deficiency (LAL-D). Diagnosis was confirmed by an enzymatic blood (e.g., dried blood spot test) or genetic test.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a specialist experienced in the treatment of inborn errors of metabolism, gastroenterologist, or lipidologist
Coverage Duration	12 months
Other Criteria	N/A



# KESIMPTA (S)

## Products Affected

- Kesimpta

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS) (Initial): Diagnosis of a relapsing form of multiple sclerosis (MS) (e.g., clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions). One of the following: 1) Failure after a trial of at least 4 weeks, contraindication, or intolerance to one of the following: A) Aubagio (teriflunomide), B) Mavenclad (cladribine), C) Plegridy (peginterferon beta-1a), D) Any one of the interferon beta-1a injections (eg, Avonex), E) Any one of the interferon beta-1b injections (eg, Betaseron, Extavia), F) Any one of the oral fumarates (eg, brand Tecfidera, generic dimethyl fumarate), G) Any one of the glatiramer acetates (eg, Copaxone, Glatopa, generic glatiramer acetate), H) Any one of the Sphingosine 1-Phosphate (S1P) receptor modulators (eg, Gilenya [fingolimod], Mayzent [siponimod], Zeposia [ozanimod]), OR 2) For continuation of prior therapy. Not used in combination with another disease-modifying therapy for MS. Not used in combination with another B-cell targeted therapy (e.g., rituximab [Rituxan], belimumab [Benlysta], ocrelizumab [Ocrevus]).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	MS (Initial, Reauth): Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	MS (Initial, Reauth): 12 months
<b>Other Criteria</b>	MS (Reauth): Documentation of positive clinical response to therapy (e.g., stability in radiologic disease activity, clinical relapse, or disease progression). Not used in combination with another disease-modifying therapy for MS. Not used in combination with another B-cell targeted therapy (e.g., rituximab [Rituxan], belimumab [Benlysta], ocrelizumab [Ocrevus]).

## KISQALI (S)

### Products Affected

- Kisqali

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: Diagnosis of advanced or metastatic breast cancer. Disease is hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative. One of the following: A) Used in combination with an aromatase inhibitor [e.g., Femara (letrozole)] OR B) Used in combination with Faslodex (fulvestrant).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy

## KISQALI-FEMARA PACK (S)

### Products Affected

- Kisqali Femara 200 Dose
- Kisqali Femara 400 Dose
- Kisqali Femara 600 Dose

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: Diagnosis of advanced or metastatic breast cancer. Disease is hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative. One of the following: A) Patient is postmenopausal OR B) Both of the following: a) Patient is pre/perimenopausal or male AND b) Treated with a Luteinizing Hormone-Releasing Hormone (LHRH) agonist (e.g. leuprolide).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## KORLYM (S)

### Products Affected

- Korlym

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cushing's syndrome (Initial): Diagnosis of endogenous Cushing's syndrome (i.e., hypercortisolism is not a result of chronic administration of high dose glucocorticoids). Diagnosis of either type 2 diabetes mellitus or diagnosis of glucose intolerance. Patient has either failed surgery or patient is not a candidate for surgery. Patient is not pregnant.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial: Prescribed by or in consultation with an endocrinologist.
<b>Coverage Duration</b>	Initial, reauth: 6 months
<b>Other Criteria</b>	Reauth: Documentation of one of the following: patient has improved glucose tolerance while on therapy or patient has stable glucose tolerance while on therapy.

## KOSELUGO (S)

### Products Affected

- Koselugo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Neurofibromatosis Type 1 (NF1): Diagnosis of NF1. Patient has plexiform neurofibromas that are both of the following: inoperable and causing significant morbidity (e.g., disfigurement, motor dysfunction, pain, airway dysfunction, visual impairment). Patient is able to swallow a capsule whole.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with one of the following: oncologist or neurologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## KUVAN (S)

### Products Affected

- Sapropterin Dihydrochloride

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Phenylketonuria (PKU) (initial): Diagnosis of PKU. Patient will have blood Phe levels measured after 1 week of therapy (new starts to therapy only) and periodically for up to 2 months of therapy to determine response.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	PKU (Init): 2 months (Reauth): 12 months
Other Criteria	PKU (reauth): Documentation of a positive clinical response to therapy. Patient will continue to have blood Phe levels measured periodically during therapy.

## KYNMOBI (S)

### Products Affected

- Kynmobi

- Kynmobi Titration Kit

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Parkinson's disease (PD) (Initial): Not used with any 5-HT3 antagonist (e.g., ondansetron, granisetron, dolasetron, palonosetron, alosetron)
Required Medical Information	Parkinson's disease (PD) (Initial): Diagnosis of PD. Patient is experiencing acute intermittent hypomobility (defined as “off” episodes characterized by muscle stiffness, slow movements, or difficulty starting movements). Used in combination with other medications for the treatment of PD (e.g., carbidopa/levodopa, pramipexole, ropinirole, etc.).
Age Restrictions	N/A
Prescriber Restrictions	PD (Initial): Prescribed by or in consultation with a neurologist.
Coverage Duration	PD (Initial, reauth): 12 months
Other Criteria	PD (Reauth): Documentation of positive clinical response to therapy.

# LEMTRADA (S)

## Products Affected

- Lemtrada

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions). One of the following: 1) Patient has not been previously treated with alemtuzumab, and failure after a trial of at least 4 weeks, contraindication, or intolerance to two of the following disease-modifying therapies for MS: A) Aubagio (teriflunomide), B) Mavenclad (cladribine), C) Plegridy (peginterferon beta-1a), D) Tysabri (natalizumab), E) Any one of the interferon beta-1a injections (eg, Avonex), F) Any one of the interferon beta-1b injections (eg, Betaseron, Extavia), G) Any one of the oral fumarates (eg, brand Tecfidera, generic dimethyl fumarate), H) Any one of the glatiramer acetate injections (eg, Copaxone, Glatopa, generic glatiramer acetate), I) Any one of the Sphingosine 1-Phosphate (S1P) receptor modulators (eg, Gilenya [fingolimod], Mayzent [siponimod], Zeposia [ozanimod]), J) Any one of the B-cell targeted therapies (eg, Ocrevus [ocrelizumab], Kesimpta [ofatumumab]), or 2) Patient has previously received treatment with alemtuzumab, and at least 12 months have or will have elapsed since the most recent treatment course with alemtuzumab. Not used in combination with another disease-modifying therapy for MS.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	MS: Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	MS: 12 months.
<b>Other Criteria</b>	N/A



# LENVIMA (S)

## Products Affected

- Lenvima 10 Mg Daily Dose
- Lenvima 12mg Daily Dose
- Lenvima 14 Mg Daily Dose
- Lenvima 18 Mg Daily Dose
- Lenvima 20 Mg Daily Dose
- Lenvima 24 Mg Daily Dose
- Lenvima 4 Mg Daily Dose
- Lenvima 8 Mg Daily Dose

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Differentiated thyroid cancer (DTC): Diagnosis of DTC. Disease is locally recurrent or metastatic. Patient has symptomatic or progressive disease. Disease is refractory to radioactive iodine treatment. Renal Cell Carcinoma (RCC): Diagnosis of advanced RCC. One of the following: 1) Both of the following: a) Used as first-line treatment and b) Used in combination with Keytruda (pembrolizumab), or 2) Both of the following: a) Treatment follows one prior anti-angiogenic therapy and b) Used in combination with everolimus. Hepatocellular Carcinoma (HCC): Diagnosis of HCC. One of the following: patient has metastatic disease, or patient has extensive liver tumor burden, or patient is inoperable by performance status or comorbidity (local disease or local disease with minimal extrahepatic disease only), or disease is unresectable. Endometrial Carcinoma (EC): Diagnosis of advanced endometrial carcinoma that is not microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR). Patient has disease progression following systemic therapy. Used in combination with Keytruda (pembrolizumab) therapy. Patient is not a candidate for curative surgery or radiation.
Age Restrictions	N/A
Prescriber Restrictions	DTC/RCC/EC: Prescribed by or in consultation with an oncologist. HCC: Prescribed by or in consultation with one of the following: oncologist, hepatologist, or gastroenterologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## LETAIRIS (S)

### Products Affected

- Ambrisentan

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
<b>Coverage Duration</b>	PAH (Initial): 6 months. PAH (Reauth): 12 months
<b>Other Criteria</b>	PAH (Reauth): Documentation of positive clinical response to therapy.

## LIDOCAINE TOPICAL (S)

### Products Affected

- Glydo
- Lidocaine OINT 5%
- Lidocaine Hcl PRSY
- Lidocaine Hcl Jelly PRSY
- Lidocaine/prilocaine CREA
- Lidocaine-prilocaine-cream Base CREA 2.5%; 2.5%
- Premium Lidocaine

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A

## LIDODERM (S)

---

### Products Affected

- Lidocaine PTCH 5%

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Post-herpetic neuralgia: Diagnosis of post-herpetic neuralgia.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

# LIVMARLI (S)

## Products Affected

- Livmarli

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Alagille syndrome (ALGS) (initial): Both of the following: a) Diagnosis of ALGS, and b) Molecular genetic testing confirms mutations in the JAG1 or NOTCH2 gene. Documentation of ONE of the following: a) Total serum bile acid greater than 3 times the upper limit of normal (ULN), b) Conjugated bilirubin greater than 1 mg/dL, c) Fat soluble vitamin deficiency otherwise unexplainable, or d) Gammaglutamyl transpeptidase (GGT) greater than 3 times the ULN. Patient is experiencing moderate to severe cholestatic pruritus. Patient has had an inadequate response to at least two of the following treatments used for the relief of pruritus: a) Ursodeoxycholic acid (e.g., Ursodiol), b) Antihistamines (e.g., diphenhydramine, hydroxyzine), c) Rifampin, or d) Bile acid sequestrants (e.g., Questran, Colestid, Welchol).
<b>Age Restrictions</b>	ALGS (initial): Patient is 1 year of age or older.
<b>Prescriber Restrictions</b>	ALGS (initial): Prescribed by or in consultation with a hepatologist.
<b>Coverage Duration</b>	ALGS (initial, reauth): 12 months.
<b>Other Criteria</b>	ALGS (reauth): Documentation of positive clinical response to therapy (e.g., reduced bile acids, reduced pruritus severity score).

# LONSURF (S)

## Products Affected

- Lonsurf

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Colorectal Cancer: Diagnosis of metastatic colorectal cancer AND trial and failure, contraindication, or intolerance to at least one component in the following: fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy (e.g., FOLFOX, FOLFIRI, FOLFOXIRI) AND trial and failure, contraindication, or intolerance to at least one anti-VEGF therapy (e.g., bevacizumab)) AND One of the following: A) patient has KRAS wild-type tumors and trial and failure, contraindication, or intolerance to at least one anti-EGFR therapy (e.g., Vectibix, Erbitux) OR Patient has KRAS mutant tumors. Gastric/Gastroesophageal Junction Adenocarcinoma: Diagnosis of metastatic gastric cancer or diagnosis of metastatic gastroesophageal junction adenocarcinoma. Trial and failure, contraindication or intolerance to at least two of the following: fluoropyrimidine-based chemotherapy (e.g. fluorouracil), Platinum-based chemotherapy (e.g., carboplatin, cisplatin, oxaliplatin), Taxane (e.g., docetaxel, paclitaxel) or irinotecan-based chemotherapy, HER2/neu-targeted therapy (e.g., trastuzumab) (if HER2 overexpression).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## LORBRENA (S)

---

### Products Affected

- Lorbrena

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC. Patient has an anaplastic lymphoma kinase (ALK)-positive tumor as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# LOTRONEX (S)

## Products Affected

- Alosetron Hydrochloride

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Severe Diarrhea-Predominant Irritable Bowel Syndrome (IBS) in Women (initial): All of the following: 1) diagnosis of severe diarrhea-predominant IBS, 2) symptoms for at least 6 months, 3) female patient, AND 4) trial and failure, contraindication, or intolerance to an antidiarrheal agent [eg, loperamide].
Age Restrictions	Initial: 18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	IBS (initial): 12 weeks. IBS (reauth): 6 mo.
Other Criteria	IBS (reauthorization): Symptoms of IBS continue to persist, AND documentation of positive clinical response to therapy.



## LUMAKRAS (S)

### Products Affected

- Lumakras

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Disease is one of the following: a) locally advanced or b) metastatic. Tumor is KRAS G12C-mutated as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient has received at least one prior systemic therapy (e.g., cisplatin/pemetrexed, atezolizumab, nivolumab, capmatinib).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## LUMIZYME (S)

---

### Products Affected

- Lumizyme

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pompe disease: Diagnosis of Pompe disease [acid alpha-glucosidase (GAA) deficiency].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

## LUPRON (S)

---

### Products Affected

- Leuprolide Acetate INJ 1MG/0.2ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prostate Cancer: Diagnosis of advanced or metastatic prostate cancer.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Prostate CA: 12 months
Other Criteria	Approve for continuation of prior therapy.

## LUPRON DEPOT (S)

### Products Affected

- Lupron Depot (1-month)
- Lupron Depot (3-month)
- Lupron Depot (4-month)
- Lupron Depot (6-month)

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prostate Cancer (7.5 mg, 22.5 mg, 30 mg, 45 mg): Diagnosis of advanced or metastatic prostate cancer. Endometriosis (3.75 mg, 11.25 mg) (initial): Diagnosis of endometriosis. One of the following: Patient has had surgical ablation to prevent recurrence, or trial and failure, contraindication, or intolerance to one NSAID (e.g., diclofenac, ibuprofen, meloxicam, naproxen) and one oral contraceptive (e.g., norethindrone-ethinyl estradiol, estradiol and norethindrone). Uterine Leiomyomata (UL) (3.75 mg, 11.25 mg): a) For use prior to surgery to reduce size of fibroids to facilitate a surgical procedure (eg, myomectomy, hysterectomy) OR b) all of the following: treatment of anemia, anemia is caused by uterine leiomyomata (fibroids), and for use prior to surgery.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Prostate CA: 12 mo. Endomet (init, reauth):6mo. UL (anemia):3 mo (fibroids):4 mo
Other Criteria	Approve for continuation of prior therapy.

# LYNPARZA TABLET (S)

## Products Affected

- Lynparza TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Ovarian cancer, advanced disease with known or suspected BRCA mutation with 3 or more prior lines of chemotherapy: Diagnosis of advanced ovarian cancer. Presence of deleterious or suspected deleterious germline BRCA-mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Trial and failure, contraindication, or intolerance to three or more prior lines of chemotherapy (e.g., paclitaxel with cisplatin). Maintenance treatment of recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of one of the following: recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer. Used for maintenance treatment in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin). Breast cancer: Diagnosis of metastatic breast cancer. Presence of a deleterious or suspected deleterious germline BRCA-mutation as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Disease is human epidermal growth factor receptor 2 (HER2)-negative. Patient has been previously treated with chemotherapy (e.g., anthracycline, taxane) in the neoadjuvant, adjuvant, or metastatic setting. One of the following: a) Disease is hormone receptor (HR) negative, or b) Disease is hormone receptor (HR)-positive and one of the following: i) patient has been treated with prior endocrine therapy or ii) patient is considered an inappropriate candidate for endocrine therapy. See Other Criteria
Age Restrictions	N/A
Prescriber Restrictions	All uses (except prostate cancer): Prescribed by or in consultation with an oncologist. Prostate cancer: Prescribed by or in consultation with an oncologist or urologist.

<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	<p>First-line maintenance treatment of BRCA-mutated advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of one of the following: advanced epithelial ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Presence of deleterious or suspected deleterious BRCA-mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient has had a complete or partial response to first-line platinum-based chemotherapy (e.g., carboplatin, cisplatin). Will be used as first-line maintenance treatment. Pancreatic adenocarcinoma: Diagnosis of metastatic pancreatic adenocarcinoma. Presence of a deleterious or suspected deleterious germline BRCA-mutation as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Disease has not progressed while receiving at least 16 weeks of a first-line platinum-based chemotherapy regimen (e.g., FOLFIRINOX, FOLFOX, etc.). First-line maintenance treatment of HRD-positive advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer in combination with bevacizumab: Diagnosis of advanced epithelial ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Cancer is associated with homologous recombination deficiency (HRD)-positive status (defined by either: a deleterious or suspected deleterious BRCA mutation, and/or genomic instability) as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient has had a complete or partial response to first-line platinum-based chemotherapy (e.g., carboplatin, cisplatin). Used in combination with bevacizumab (e.g., Avastin, Mvasi). Will be used as first-line maintenance treatment. Prostate cancer: Diagnosis of metastatic castration-resistant prostate cancer. Presence of deleterious or suspected deleterious homologous recombination repair (HRR) gene mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Disease has progressed following prior treatment with one of the following: a) enzalutamide (Xtandi) or b) abiraterone (e.g., Zytiga, Yonsa). All indications: Approve for continuation of prior therapy.</p>

## MAKENA (S)

### Products Affected

- Makena INJ 275MG/1.1ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Preterm birth prophylaxis: Patient had a previous singleton (single offspring) spontaneous preterm birth. Patient is having a singleton pregnancy. Therapy will be started between 16 weeks, 0 days and 20 weeks, 6 days of gestation. Therapy will be continued until week 37 (through 36 weeks, 6 days) of gestation or delivery, whichever occurs first.
Age Restrictions	N/A
Prescriber Restrictions	Preterm birth prophylaxis: Prescribed by or in consultation with a specialist in obstetrics and gynecology
Coverage Duration	Preterm birth prophylaxis: 21 weeks
Other Criteria	N/A

## MARINOL (S)

### Products Affected

- Dronabinol

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Nausea and Vomiting Associated with Cancer Chemotherapy (CINV): Patient is receiving cancer chemotherapy. Trial and failure, contraindication, or intolerance to one 5HT-3 receptor antagonist (eg, Anzemet [dolasetron], Kytril [granisetron], or Zofran [ondansetron]). Trial and failure, contraindication, or intolerance to one of the following: Compazine (prochlorperazine), Decadron (dexamethasone), Haldol (haloperidol), Zyprexa (olanzapine). AIDS anorexia: Diagnosis of anorexia with weight loss in patients with AIDS. Patient is on antiretroviral therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	CINV: 6 months. AIDS anorexia: 3 months.
<b>Other Criteria</b>	Subject to Part B vs. Part D review. CINV: Approve for continuation of therapy for treatment covered under Part B when patient is receiving cancer chemotherapy.



## MAVYRET (S)

### Products Affected

- Mavyret

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guideline. All patients: Diagnosis of chronic hepatitis C, patient is without decompensated liver disease (defined as Child-Pugh Class B or C), and not used in combination with another HCV direct acting antiviral agent [e.g., Harvoni, Zepatier].
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine.
Coverage Duration	8 to 16 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline.
Other Criteria	N/A

# MEKINIST (S)

## Products Affected

- Mekinist

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Melanoma: Diagnosis of unresectable or metastatic melanoma AND cancer is BRAF V600E or V600K mutant type as detected by a U.S. Food and Drug Administration (FDA)-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Adjuvant Treatment for Melanoma: Diagnosis of melanoma. Cancer is BRAF V600E or V600K mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Involvement of lymph nodes following complete resection. Used as adjunctive therapy. Medication is used in combination with Tafenlar (dabrafenib). Non-small Cell Lung Cancer (NSCLC): All of the following: diagnosis of metastatic non-small cell lung cancer AND cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) AND medication is used in combination with Tafenlar (dabrafenib). Anaplastic Thyroid Cancer (ATC): Diagnosis of locally advanced or metastatic ATC. Cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Cancer may not be treated with standard locoregional treatment options. Medication is used in combination with Tafenlar (dabrafenib).</p>
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## MEKTOVI (S)

### Products Affected

- Mektovi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Melanoma: Diagnosis of unresectable melanoma or metastatic melanoma. Cancer is BRAF V600E or V600K mutant type (MT) as detected by a U.S. Food and Drug Administration (FDA)-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Used in combination with Braftovi (encorafenib).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy

## MIGRANAL (S)

### Products Affected

- Dihydroergotamine Mesylate NASAL SOLN

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial: Diagnosis of migraine headaches with or without aura. Will be used for the acute treatment of migraine. One of the following: Trial and failure or intolerance to one triptan (e.g., eletriptan, rizatriptan, sumatriptan) or contraindication to all triptans.
Age Restrictions	N/A
Prescriber Restrictions	Initial, Reauth: Prescribed by or in consultation with a neurologist, headache specialist, or pain specialist.
Coverage Duration	Initial: 3 months. Reauth: 12 months.
Other Criteria	Reauth: Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea).

## MONJUVI (S)

### Products Affected

- Monjuvi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diffuse Large B-cell Lymphoma (DLBCL): Diagnosis of DLBCL. Disease is one of the following: relapsed or refractory. Used in combination with lenalidomide. Patient is not eligible for autologous stem cell transplant (ASCT).
Age Restrictions	N/A
Prescriber Restrictions	DLBCL: Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## MS INTERFERONS (PREFERRED) (S)

---

### Products Affected

- Betaseron

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple Sclerosis (MS) (initial): Diagnosis of a relapsing form of MS (eg, clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions).
Age Restrictions	N/A
Prescriber Restrictions	MS (initial, reauth): Prescribed by or in consultation with a neurologist
Coverage Duration	MS (initial, reauth): 12 months
Other Criteria	MS (reauth): Documentation of positive clinical response to therapy (e.g., stability in radiologic disease activity, clinical relapses, disease progression).

# MVASI (S)

## Products Affected

- Mvasi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Non-Small Cell Lung Cancer: Excluded if squamous cell histology.
Required Medical Information	<p>Colorectal Cancer: Diagnosis of metastatic colorectal cancer. One of the following: 1) Both of the following: a) used as first- or second-line treatment and b) used in combination with an intravenous 5-fluorouracil-based chemotherapy, OR 2) All of the following: a) used as second-line treatment, b) used in combination with fluoropyrimidine-irinotecan-based chemotherapy or fluoropyrimidine-oxaliplatin-based chemotherapy, and c) patient has progressed on a first-line bevacizumab-containing regimen.</p> <p>Non-Small Cell Lung Cancer (NSCLC): Diagnosis of NSCLC. Disease is unresectable, locally advanced, recurrent, or metastatic. Used as first-line treatment. Used in combination with paclitaxel and carboplatin.</p> <p>Renal Cell Cancer: Diagnosis of metastatic renal cell cancer. Used in combination with interferon-alpha.</p> <p>Cervical Cancer: Diagnosis of carcinoma of the cervix. Disease is persistent, recurrent, or metastatic. Used in combination with one of the following: a) paclitaxel and cisplatin or b) paclitaxel and topotecan.</p> <p>Glioblastoma: Diagnosis of recurrent glioblastoma.</p> <p>Epithelial Ovarian, Fallopian Tube, or Primary Peritoneal Cancer: Diagnosis of epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. One of the following: 1) All of the following: a) disease is stage 3 or 4, b) patient has been treated with bevacizumab as a single agent, c) treatment is following surgical resection, and d) used in combination with carboplatin and paclitaxel, OR 2) All of the following: a) disease is platinum-resistant recurrent, b) patient has received no more than 2 prior chemotherapy regimens, and c) used in combination with paclitaxel, pegylated liposomal doxorubicin, or topotecan, OR 3) All of the following: a) disease is platinum-sensitive recurrent, b) patient has been treated with bevacizumab as a single agent, and c) used in combination with one of the following: i) carboplatin and paclitaxel or ii) carboplatin and gemcitabine.</p>
Age Restrictions	N/A

<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Hepatocellular Carcinoma (HCC): Diagnosis of hepatocellular carcinoma. Disease is unresectable or metastatic. Used in combination with Tecentriq (atezolizumab). Patient has not received prior systemic therapy. Approve for continuation of prior therapy.



## MYALEPT (S)

### Products Affected

- Myalept

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Lipodystrophy (initial): Diagnosis of congenital or acquired generalized lipodystrophy AND one of the following: 1) Diabetes mellitus or insulin resistance despite optimized insulin therapy at maximum tolerated doses OR 2) Hypertriglyceridemia despite optimized therapy with at least two triglyceride-lowering agents from different classes (e.g., fibrates, statins).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial: Prescribed by or in consultation with an endocrinologist
<b>Coverage Duration</b>	Initial, reauth: 12 months
<b>Other Criteria</b>	Lipodystrophy (reauth): Patient has experienced an objective response to therapy, such as A) Sustained reduction in hemoglobin A1c (HbA1c) level from baseline OR B) Sustained reduction in triglyceride (TG) levels from baseline.

## NAGLAZYME (S)

---

### Products Affected

- Naglazyme

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Mucopolysaccharidosis (MPS VI): Diagnosis of MPS VI (Maroteaux-Lamy Syndrome).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	MPS VI: 12 months
Other Criteria	N/A

## NATPARA (S)

### Products Affected

- Natpara

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Hypocalcemia (Initial): Diagnosis of hypocalcemia due to chronic hypoparathyroidism. Not used in the setting of acute post-surgical hypoparathyroidism. Patient does not have a known calcium-sensing receptor mutation. Patient has a documented parathyroid hormone concentration that is inappropriately low for the level of calcium, recorded on at least two occasions within the previous 12 months. Patient has normal thyroid-stimulating hormone concentrations if not on thyroid hormone replacement therapy (or if on therapy, the dose had to have been stable for greater than or equal to 3 months). Patient has normal magnesium and serum 25-hydroxyvitamin D concentrations. Will be used as an adjunct treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Hypocalcemia (initial): Prescribed by or in consultation with an endocrinologist.
<b>Coverage Duration</b>	Initial: 6 months. Reauth: 12 months
<b>Other Criteria</b>	Hypocalcemia (Reauth): One of the following: A) Patient has achieved and maintained serum calcium levels in the ideal range (7.5 - 10.6 mg/dL), OR B) Patient has experienced a 50% or greater reduction in oral calcium intake, OR C) Patient has experienced a 50% or greater reduction in oral vitamin D intake.

## NERLYNX (S)

### Products Affected

- Nerlynx

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Early Stage Breast cancer: Diagnosis (dx) of early stage breast cancer. Disease is human epidermal growth factor receptor 2 (HER2)-positive. Patient has received adjuvant trastuzumab-based therapy. Advanced or Metastatic Breast Cancer: Dx of advanced or metastatic breast cancer. Disease is human epidermal growth factor receptor 2 (HER2)-positive. Patient has received two or more prior anti-HER2 based regimens (e.g., trastuzumab + pertuzumab + docetaxel, ado-trastuzumab emtansine, etc.). Used in combination with capecitabine.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# NEULASTA (S)

## Products Affected

- Neulasta

- Neulasta Onpro Kit

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Febrile neutropenia (FN) prophylaxis: Patient will be receiving prophylaxis for FN due to one of the following: 1) Patient is receiving National Cancer Institute's Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer, 2) patient is receiving a dose-dense chemotherapy regimen for which the incidence of FN is unknown, 3) patient is receiving chemotherapy regimen(s) associated with greater than 20% incidence of FN, 4) both of the following: a) patient is receiving chemotherapy regimen(s) associated with 10-20% incidence of FN, AND b) patient has one or more risk factors associated with chemotherapy-induced infection, FN, or neutropenia, OR 5) Both of the following: a) patient is receiving myelosuppressive anticancer drugs associated with neutropenia, AND b) patient has a history of FN or dose-limiting event during a previous course of chemotherapy (secondary prophylaxis). Acute radiation syndrome (ARS): Patient was/will be acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of ARS). Treatment of FN: Patient has received or is receiving myelosuppressive anticancer drugs associated with neutropenia. Diagnosis of FN. Patient is at high risk for infection-associated complications.
Age Restrictions	N/A
Prescriber Restrictions	All uses: Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	ARS: 1 mo. FN (prophylaxis, treatment): 3 mo or duration of tx.
Other Criteria	N/A

# NEXAVAR (S)

## Products Affected

- Nexavar

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Renal cell carcinoma (RCC): Diagnosis of RCC. Hepatocellular carcinoma (HCC): Diagnosis of HCC. One of the following: patient has metastatic disease, or patient has extensive liver tumor burden, or patient is inoperable by performance status or comorbidity (local disease or local disease with minimal extrahepatic disease only), or disease is unresectable. Differentiated thyroid carcinoma (DTC): Diagnosis of DTC (ie, follicular carcinoma, Hurthle cell carcinoma, or papillary carcinoma). One of the following: locally recurrent disease, metastatic disease, or unresectable disease. One of the following: patient has symptomatic disease or patient has progressive disease. Disease is refractory to radioactive iodine (RAI) treatment. Medullary thyroid carcinoma (MTC): Diagnosis of MTC. One of the following: 1) Disease is progressive or 2) Disease is symptomatic with distant metastases. Trial and failure, contraindication, or intolerance to Caprelsa (vandetanib) or Cometriq (cabozantinib).
Age Restrictions	N/A
Prescriber Restrictions	DTC, MTC: Prescribed by or in consultation with an oncologist. RCC: Prescribed by or in consultation with an oncologist or nephrologist. HCC: Prescribed by or in consultation with an oncologist, hepatologist, or gastroenterologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## NINLARO (S)

---

### Products Affected

- Ninlaro

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple myeloma: Diagnosis of multiple myeloma. Used in combination with Revlimid (lenalidomide) and dexamethasone. Patient has received at least one prior therapy for multiple myeloma [eg, Revlimid (lenalidomide), Thalomid (thalidomide), Velcade (bortezomib)].
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist/oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## NORTHERA (S)

### Products Affected

- Droxidopa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Neurogenic orthostatic hypotension (NOH) (init): Diagnosis of symptomatic NOH. NOH is caused by one of the following conditions: primary autonomic failure (eg, Parkinson's disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, non-diabetic autonomic neuropathy. Trial and failure, contraindication, or intolerance to one of the following agents: fludrocortisone acetate, midodrine.
Age Restrictions	N/A
Prescriber Restrictions	NOH (init): Prescribed by or in consultation with a cardiologist, neurologist, or nephrologist
Coverage Duration	NOH (init): 1 month (reauth): 12 months
Other Criteria	NOH (reauth): Documentation of positive clinical response to therapy.



## NUBEQA (S)

### Products Affected

- Nubeqa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-metastatic castration-resistant or castration-recurrent prostate cancer (nmCRPC): Diagnosis of non-metastatic castration-resistant (chemical or surgical) or castration-recurrent prostate cancer.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## NUEDEXTA (S)

### Products Affected

- Nuedexta

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Pseudobulbar affect (PBA) (initial): Diagnosis of PBA. Patient does not have any of the following contraindications: a) Concomitant use with other drugs containing quinidine, quinine, or mefloquine, b) History of Nuedexta, quinine, mefloquine or quinidine-induced thrombocytopenia, hepatitis, bone marrow depression, or lupus-like syndrome, c) Known hypersensitivity to dextromethorphan (e.g., rash, hives), d) Taking monoamine oxidase inhibitors (MAOIs) (e.g., phenelzine, selegiline, tranlycypromine) or have taken MAOIs within the preceding 14 days, e) Has prolonged QT interval, congenital long QT syndrome or a history suggestive of torsades de pointes, or has heart failure, f) Receiving drugs that both prolong QT interval and are metabolized by CYP2D6 (e.g., thioridazine, pimozide), g) Has complete atrioventricular (AV) block without implanted pacemakers, or at high risk of complete AV block.</p> <p>PBA (reauth): Documentation of clinical benefit from ongoing therapy as demonstrated by a decrease in inappropriate laughing or crying episodes.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PBA (initial): Prescribed by or in consultation with one of the following specialists: neurologist, psychiatrist.
<b>Coverage Duration</b>	PBA (initial/reauth): 12 months
<b>Other Criteria</b>	N/A

## NUPLAZID (S)

---

### Products Affected

- Nuplazid

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Parkinson's disease psychosis: Diagnosis of Parkinson's disease. Patient has at least one of the following: hallucinations or delusions.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# NUVIGIL (S)

## Products Affected

- Armodafinil

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Obstructive sleep apnea (OSA) (Initial): Diagnosis (dx) of OSA defined by one of the following: a) 15 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), or b) both of the following: 5 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), AND 1 of the following symptoms: unintentional sleep episodes during wakefulness, daytime sleepiness, unrefreshing sleep, fatigue, insomnia, waking up breath holding/gasping/choking, loud snoring, or breathing interruptions during sleep. Shift-work disorder (SWD) (Initial):Dx of SWD confirmed by one of the following: 1) symptoms of excessive sleepiness or insomnia for at least 3 months, which is associated with a work period (usually night work) that occurs during the normal sleep period, OR 2) A sleep study demonstrating loss of a normal sleep-wake pattern (ie, disturbed chronobiologic rhythmicity). Confirmation that no other medical conditions or medications are causing the symptoms of excessive sleepiness or insomnia. Narcolepsy (initial): Dx of narcolepsy as confirmed by sleep study (unless prescriber provides justification confirming that a sleep study is not feasible).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	OSA, SWD: Initial, Reauth: 6 mo. Narcolepsy: Initial, Reauth: 12 mo
<b>Other Criteria</b>	OSA, Narcolepsy (Reauth): Documentation of positive clinical response to armodafinil therapy. SWD (Reauth): Documentation of positive clinical response to armodafinil therapy.

## ODOMZO (S)

---

### Products Affected

- Odomzo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Basal cell carcinoma: Diagnosis of locally advanced basal cell carcinoma AND One of the following: 1) Cancer has recurred following surgery or radiation therapy or 2) Patient is not a candidate for surgery or radiation therapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## OFEV (S)

### Products Affected

- Ofev

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Idiopathic pulmonary fibrosis (IPF) (initial): Diagnosis of IPF as documented by all of the following: a) exclusion of other known causes of interstitial lung disease (ILD) (eg, domestic and occupational environmental exposures, connective tissue disease, drug toxicity) AND b) one of the following: i) in patients not subjected to surgical lung biopsy, the presence of a usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) revealing IPF or probable IPF, OR ii) in patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern revealing IPF or probable IPF. Systemic sclerosis-associated interstitial lung disease (SSc-ILD) (initial): Diagnosis of SSc-ILD as documented by all of the following: a) exclusion of other known causes of ILD (eg, domestic and occupational environmental exposures, connective tissue disease, drug toxicity) AND b) One of the following: i) In patients not subjected to surgical lung biopsy, the presence of idiopathic interstitial pneumonia (eg, fibrotic nonspecific interstitial pneumonia [NSIP], usual interstitial pneumonia [UIP] and centrilobular fibrosis) pattern on HRCT revealing SSc-ILD or probable SSc-ILD, OR ii) in patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern revealing SSc-ILD or probable SSc-ILD. Chronic Fibrosing Interstitial Lung Diseases (ILDs) with a Progressive Phenotype (initial): 1) diagnosis of chronic fibrosing interstitial lung disease, AND 2) patient has a high-resolution computed tomography (HRCT) showing at least 10% of lung volume with fibrotic features, AND 3) disease has a progressive phenotype as observed by one of the following: decline of forced vital capacity (FVC), worsening of respiratory symptoms, or increased extent of fibrosis seen on imaging.</p>
Age Restrictions	N/A
Prescriber Restrictions	IPF, SSc-ILD, Chronic Fibrosing ILDs with a Progressive Phenotype (initial): Prescribed by or in consultation with a pulmonologist

<b>Coverage Duration</b>	Initial, reauth: 12 months
<b>Other Criteria</b>	IPF, SSc-ILD, Chronic Fibrosing ILDs with a Progressive Phenotype (reauth): Documentation of positive clinical response to therapy.

## ONUREG (S)

### Products Affected

- Onureg

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute Myeloid Leukemia (AML): Diagnosis of acute myeloid leukemia (AML). Patient has received previous treatment with an intensive induction chemotherapy regimen (e.g., cytarabine + daunorubicin, cytarabine + idarubicin, etc.). Patient has achieved one of the following: a) first complete remission (CR) or b) complete remission with incomplete blood count recovery (CRi). Patient is not able to complete intensive curative therapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.



## OPSUMIT (S)

### Products Affected

- Opsumit

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
Age Restrictions	N/A
Prescriber Restrictions	PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
Coverage Duration	PAH: Initial: 6 months. Reauth: 12 months.
Other Criteria	PAH (Reauth): Documentation of positive clinical response to therapy.

# ORENCIA IV (S)

## Products Affected

- Orencia INJ 250MG

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: a) Trial and failure, contraindication, or intolerance (TF/C/I) to two of the following: Enbrel (etanercept), Humira (adalimumab), Rinvoq (upadacitinib), or Xeljanz/Xeljanz XR (tofacitinib), or b) attestation demonstrating a trial may be inappropriate, OR c) For continuation of prior Orencia therapy. Juvenile Idiopathic Arthritis (JIA) (Initial): Diagnosis of moderately to severely active polyarticular JIA. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. JIA, PsA (Initial): One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RA, JIA (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist.
<b>Coverage Duration</b>	All uses (initial, reauth): 12 months
<b>Other Criteria</b>	All indications (Reauth): Documentation of positive clinical response to therapy.

# ORENCIA SC (S)

## Products Affected

- Orenzia INJ 125MG/ML, 50MG/0.4ML, 87.5MG/0.7ML

- Orenzia Clickject

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: a) Either a trial and failure, contraindication, or intolerance (TF/C/I) to two of the following: Enbrel (etanercept), Humira (adalimumab), Rinvoq (upadacitinib), Xeljanz/Xeljanz XR (tofacitinib), or attestation demonstrating a trial may be inappropriate, OR b) For continuation of prior therapy. Juvenile Idiopathic Arthritis (JIA) (Initial): Diagnosis of moderately to severely active polyarticular JIA. One of the following: a) Either a trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR b) for continuation of prior therapy. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: a) Either a trial and failure, contraindication, or intolerance to two of the following, or attestation demonstrating a trial may be inappropriate: Enbrel (etanercept), Humira (adalimumab), Skyrizi (risankizumab-rzaa), Rinvoq (upadacitinib), Xeljanz/XR (tofacitinib/ER), OR b) for continuation of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	RA, JIA (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist.
Coverage Duration	All uses (initial, reauth): 12 months
Other Criteria	All indications (Reauth): Documentation of positive clinical response to therapy.

## ORGOVYX (S)

### Products Affected

- Orgovyx

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prostate Cancer: Diagnosis of advanced prostate cancer. Disease is one of the following: 1) Evidence of biochemical or clinical relapse following local primary intervention with curative intent or 2) Newly diagnosed androgen-sensitive metastatic disease or 3) Advanced localized disease unlikely to be cured by local primary intervention with curative intent.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an urologist or oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# ORKAMBI (S)

## Products Affected

- Orkambi TABS

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystic Fibrosis (CF) (Initial): Diagnosis of CF. Patient is homozygous for the F508del mutation in the CF transmembrane conductance regulator (CFTR) gene as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	CF (Initial): Patient is 6 years of age or older
<b>Prescriber Restrictions</b>	CF (initial): Prescribed by or in consultation with a specialist affiliated with a CF care center or pulmonologist
<b>Coverage Duration</b>	CF (initial, reauth): 12 months
<b>Other Criteria</b>	CF (Reauth): Patient is benefiting from treatment (i.e. improvement in lung function [forced expiratory volume in one second (FEV1)], decreased number of pulmonary exacerbations).

# ORKAMBI GRANULES (S)

## Products Affected

- Orkambi PACK

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystic Fibrosis (CF) (Initial): Diagnosis of CF. Patient is homozygous for the F508del mutation in the CF transmembrane conductance regulator (CFTR) gene as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). One of the following: A) Patient is 2 through 5 years of age, OR B) Both of the following: Patient is 6 years of age or greater AND Patient is unable to swallow oral tablets.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CF (Initial): Prescribed by or in consultation with a specialist affiliated with a CF care center or pulmonologist
<b>Coverage Duration</b>	CF (initial, reauth): 12 months
<b>Other Criteria</b>	CF (Reauth): Patient is benefiting from treatment (i.e., improvement in lung function [forced expiratory volume in one second (FEV1)], decreased number of pulmonary exacerbations). One of the following: A) Patient is 2 through 5 years of age, OR B) Both of the following: Patient is 6 years of age or greater AND Patient is unable to swallow oral tablets.

## OSPHERA (S)

---

### Products Affected

- Osphena

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Dyspareunia (initial): Diagnosis of moderate to severe dyspareunia due to vulvar and vaginal atrophy associated with menopause. Vaginal dryness (initial): Diagnosis of moderate to severe vaginal dryness due to vulvar and vaginal atrophy associated with menopause.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	All uses (Initial, reauth): 12 months
Other Criteria	Dyspareunia, Vaginal dryness (reauth): Documentation of positive clinical response to therapy.

## OXANDRIN (S)

### Products Affected

- Oxandrolone TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Promote weight gain (initial): Used as adjunctive therapy to promote weight gain AND Diagnosis of one of the following: Extensive surgery, Chronic infections, Severe trauma, Failure to gain or maintain at least 90% of ideal body weight without definite pathophysiologic reasons AND a nutritional consult was performed. Counterbalance protein catabolism (initial): Used to counterbalance protein catabolism associated with chronic corticosteroid administration. Bone pain: Diagnosis of bone pain associated with osteoporosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	bone pain: 1 month. Others (initial, reauth): 3 months
Other Criteria	All diagnoses except bone pain (reauth): Documentation of a positive clinical response to therapy as evidenced by an improvement in weight gain or increase in lean body mass.



## OXBRYTA (S)

### Products Affected

- Oxbryta TBSO

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Initial: Diagnosis of Sickle Cell Disease. Documentation of hemoglobin level that does not exceed 10.5 g/dL prior to therapy initiation. Trial and failure or inadequate response, contraindication, or intolerance to hydroxyurea.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial: Prescribed by or in consultation with one of the following: 1) Hematologist/Oncologist or 2) Specialist w/ expertise in the diagnosis and management of sickle cell disease.
<b>Coverage Duration</b>	Initial, Reauth: 12 months.
<b>Other Criteria</b>	Reauth: Documentation of positive clinical response to therapy (e.g., an increase in hemoglobin level of 1 g/dL or greater from baseline, decreased annualized incidence rate of vaso-occlusive crises [VOCs]). Documentation of hemoglobin level that does not exceed 10.5 g/dL.

## PEGASYS (S)

### Products Affected

- Pegasis
- Pegasis Proclick INJ 180MCG/0.5ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic hepatitis B: Diagnosis of chronic hepatitis B infection, and patient is without decompensated liver disease. Chronic Hepatitis C: Criteria will be applied consistent with current AASLD-IDSA guidance.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	HepB: 48 wks. HepC: Initial: 28 wks. Reauth: 20 wks.
Other Criteria	N/A

## PEMAZYRE (S)

---

### Products Affected

- Pemazyre

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of cholangiocarcinoma. Disease is one of the following: unresectable locally advanced or metastatic. Disease has presence of a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement. Patient has been previously treated.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hepatologist, gastroenterologist, or oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## PENNSAID (S)

### Products Affected

- Diclofenac Sodium EXTERNAL SOLN 1.5%

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Osteoarthritis of the knees (initial): Diagnosis of osteoarthritis of the knees. Patient meets one of the following: 1) Treatment failure with at least two prescription strength topical or oral non-steroidal anti-inflammatory drugs (NSAIDs) (e.g., diclofenac, ibuprofen, meloxicam, naproxen) OR 2) History of peptic ulcer disease/gastrointestinal bleed OR 3) Patient is older than 65 years of age with one additional risk factor for gastrointestinal adverse events (e.g. use of anticoagulants, chronic corticosteroids).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial, reauth: 12 months
Other Criteria	Osteoarthritis of the knees (reauth): Documentation of positive clinical response to therapy (e.g., improvement in pain symptoms of osteoarthritis).

## PHESGO (S)

### Products Affected

- Phesgo

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Early Breast Cancer (EBC): Diagnosis of breast cancer. Used in combination with chemotherapy. Disease is human epidermal growth factor receptor 2 (HER2)-positive. One of the following: a) Used for neoadjuvant treatment and disease is one of the following: locally advanced, inflammatory, or early stage breast cancer (either greater than 2 cm in diameter or node positive), OR b) Used for adjuvant treatment and disease is early breast cancer at high risk of recurrence. Metastatic Breast Cancer (MBC): Diagnosis of breast cancer. Used in combination with docetaxel. Disease is human epidermal growth factor receptor 2 (HER2)-positive metastatic breast cancer. Patient has not received prior anti-HER2 therapy or chemotherapy for metastatic disease.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## PIQRAY (S)

### Products Affected

- Piqray 200mg Daily Dose
- Piqray 250mg Daily Dose
- Piqray 300mg Daily Dose

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast Cancer (BC): Diagnosis of advanced or metastatic BC. Disease is hormone receptor (HR)-positive, and human epidermal growth factor receptor 2 (HER2)-negative. Cancer is PIK3CA-mutated as detected by an FDA-approved test (therascreen PIK3CA RGQ PCR Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient is a postmenopausal woman or male. Used in combination with fulvestrant. Disease has progressed on or after an endocrine-based regimen.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## POLIVY (S)

### Products Affected

- Polivy

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diffuse large B-cell lymphoma (DLBCL): Diagnosis of diffuse large B-cell lymphoma (DLBCL). Disease is relapsed or refractory. Used in combination with bendamustine and a rituximab product. Patient has received at least two prior therapies for DLBCL (e.g., RCHOP [rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone], HSCT [hematopoietic stem cell transplantation], CAR T [chimeric antigen receptor T-cell] therapy, RCEPP [rituximab, cyclophosphamide, etoposide, prednisone, procarbazine], GemOx [gemcitabine, oxaliplatin] with or without rituximab).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

# POMALYST (S)

## Products Affected

- Pomalyst

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Myeloma (MM): Diagnosis of MM. Used in combination with dexamethasone. Patient has received two prior therapies, including Revlimid (lenalidomide) and a proteasome inhibitor [eg, Velcade (bortezomib) or Kyprolis (carfilzomib)] or has a contraindication or intolerance to Revlimid and proteasome inhibitors. Patient has experienced disease progression on or within 60 days of completion of last therapy. Kaposi sarcoma (KS): One of the following: 1) Both of the following: a) Diagnosis of AIDS-related KS and b) Patient has failed highly active antiretroviral therapy (HAART) [e.g., Biktarvy (bictegravir/emtricitabine/tenofovir alafenamide), Dovato (dolutegravir/lamivudine), Triumeq (dolutegravir/abacavir/lamivudine)], OR 2) Both of the following: a) Diagnosis of KS and b) Patient is HIV-negative.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All indications: Prescribed by or in consultation with an oncologist or hematologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.



# POSACONAZOLE (S)

## Products Affected

- Noxafil SUSP
- Posaconazole Dr

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prophylaxis of systemic fungal infections (SFI): Used as prophylaxis of invasive fungal infections caused by Aspergillus or Candida for one of the following conditions: 1) Patient is at high risk of infections due to severe immunosuppression from hematopoietic stem cell transplant (HSCT) with graft-versus-host disease (GVHD) or hematologic malignancies with prolonged neutropenia from chemotherapy OR 2) patient has a prior fungal infection requiring secondary prophylaxis. Treatment (Tx) of SFI: Used as treatment of systemic fungal infections caused by Aspergillus. Oropharyngeal Candidiasis (OPC): Diagnosis of OPC. One of the following: 1) Candidiasis is refractory or resistant to treatment with fluconazole OR 2) Trial and failure, contraindication, or intolerance to fluconazole.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Prophylaxis of SFI: 6 months. Tx of SFI: 3 months. OPC: 1 month.
Other Criteria	N/A

# PROMACTA (S)

## Products Affected

- Promacta

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Immune (idiopathic) thrombocytopenic purpura (ITP) (initial): Diagnosis of one of the following: relapsed/refractory ITP, persistent ITP, or chronic ITP. Baseline platelet count is less than 30,000/mcL. Patient's degree of thrombocytopenia and clinical condition increase the risk of bleeding. Trial and failure, intolerance, contraindication to corticosteroids (e.g., prednisone, methylprednisolone), immunoglobulins [e.g., Gammagard, immune globulin (human)], or splenectomy. Chronic hepatitis C (initial): Diagnosis of chronic hepatitis C-associated thrombocytopenia. One of the following: 1) Planning to initiate and maintain interferon-based treatment, or 2) currently receiving interferon-based treatment. First-line for severe aplastic anemia (SAA): Diagnosis of SAA. Used for first-line treatment (i.e., patient has not received prior immunosuppressive therapy). Used in combination with standard immunosuppressive therapy. Patient meets at least two of the following: 1) absolute neutrophil count less than 500/mcL, 2) platelet count less than 20,000/mcL, 3) absolute reticulocyte count less than 60,000/mcL. Refractory SAA (initial): Diagnosis of refractory severe aplastic anemia. Patient has a platelet count less than 30,000/mcL. Insufficient response to immunosuppressive therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	ITP and SAA: Prescribed by or in consultation with a hematologist/oncologist. Chronic hepatitis C associated thrombocytopenia: Prescribed by or in consultation with a hematologist/oncologist, gastroenterologist, hepatologist, infectious disease specialist, or HIV specialist certified through the American Academy of HIV Medicine.
<b>Coverage Duration</b>	ITP(init, reauth): 12mo. HepC: 3mo(init), 12mo(reauth). 1st line SAA: 6mo. Refract SAA: 16wk-init, 12mo-reauth

<b>Other Criteria</b>	<p>ITP (reauth): Documentation of positive clinical response to therapy as evidenced by an increase in platelet count to a level sufficient to avoid clinically important bleeding. Hepatitis C (reauth): One of the following:</p> <p>1) For patients that started treatment with eltrombopag prior to initiation of treatment with interferon, eltrombopag will be approved when both of the following are met: a) Currently on antiviral interferon therapy for treatment of chronic hepatitis C and b) Documentation that the patient reached a threshold platelet count that allows initiation of antiviral interferon therapy with eltrombopag treatment by week 9, OR</p> <p>2) For patients that started treatment with eltrombopag while on concomitant treatment with interferon, eltrombopag will be approved based on the following: Currently on antiviral interferon therapy for treatment of chronic hepatitis C. Refractory SAA (reauth): Documentation of positive clinical response to therapy as evidenced by an increase in platelet count.</p>
-----------------------	---

# PROVIGIL (S)

## Products Affected

- Modafinil

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Obstructive sleep apnea (OSA) (Initial): Diagnosis (dx) of OSA defined by one of the following: 15 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), or both of the following: 5 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), and 1 of the following symptoms: unintentional sleep episodes during wakefulness, daytime sleepiness, unrefreshing sleep, fatigue, insomnia, waking up breath holding/gasping/choking, loud snoring, or breathing interruptions during sleep. Shift-work disorder (SWD) (Initial): Dx of SWD confirmed by one of the following: 1) Symptoms of excessive sleepiness or insomnia for at least 3 months, which is associated with a work period (usually night work) that occurs during the normal sleep period, OR 2) A sleep study demonstrating loss of a normal sleep-wake pattern (ie, disturbed chronobiologic rhythmicity). Confirmation that no other medical conditions or medications are causing the symptoms of excessive sleepiness or insomnia. Narcolepsy (initial): Dx of narcolepsy as confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible). MS Fatigue (initial): Dx of multiple sclerosis (MS). Patient is experiencing fatigue. Depression (initial): Treatment-resistant depression defined as diagnosis of major depressive disorder (MDD) or bipolar depression, AND trial and failure, contraindication, or intolerance to at least two antidepressants from different classes (eg, SSRIs, SNRIs, bupropion). Used as adjunctive therapy. Idiopathic Hypersomnia (Initial): Diagnosis of idiopathic hypersomnia as confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible).
Age Restrictions	N/A

<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Narcolepsy: Init, Reauth: 12 mo. All other indications: Init, Reauth: 6 mo.
<b>Other Criteria</b>	OSA, Narcolepsy, Idiopathic Hypersomnia (Reauth): Documentation of positive clinical response to modafinil therapy. SWD (Reauth): Documentation of positive clinical response to modafinil therapy. MS Fatigue (reauth): Patient is experiencing relief of fatigue with modafinil therapy. Depression (reauth): Documentation of positive clinical response to modafinil therapy. Used as adjunctive therapy.

## QINLOCK (S)

---

### Products Affected

- Qinlock

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Gastrointestinal Stromal Tumor (GIST): Diagnosis of gastrointestinal stromal tumor (GIST). Disease is advanced. Patient has received prior treatment with three or more kinase inhibitors (e.g., sunitinib, regorafenib), one of which must include imatinib.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## QUALAQUIN (S)

### Products Affected

- Quinine Sulfate CAPS 324MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Excluded if used solely for the treatment or prevention of nocturnal leg cramps.
Required Medical Information	Malaria: Diagnosis of uncomplicated malaria. One of the following: 1) Treatment in areas of chloroquine-sensitive malaria, and trial and failure, contraindication, or intolerance to chloroquine or hydroxychloroquine, OR 2) Treatment in areas of chloroquine-resistant malaria.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	7 days
Other Criteria	N/A

# REPATHA (S)

## Products Affected

- Repatha

- Repatha Pushtronex System
- Repatha Sureclick

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>HeFH/ASCVD/Primary HLD (init): One of the following dx: A)HeFH as confirmed by one of the following: 1)Both of the following: a)Untreated/pre-treatment LDL greater than 190 mg/dL in an adult, AND b)One of the following: i) Family hx of tendinous xanthomas and/or arcus cornealis in 1st degree relative, or 2nd degree relative, ii)Hx of myocardial infarction (MI) in 1st-degree relative less than 60 years of age, iii)Family hx of MI in 2nd-degree relative less than 50 years of age, iv)Family hx of LDL-C greater than 190 mg/dL in 1st- or 2nd-degree relative, v)Family hx of FH in 1st- or 2nd-degree relative, or 2)Untreated/pre-treatment LDL-C greater than 190 mg/dL in an adult AND submission of MR (eg chart notes, lab values) documenting one of the following: presence of tendinous xanthoma in pt, arcus cornealis before age 45, or functional mutation in the LDL receptor, ApoB, or PCSK9 gene. OR B)ASCVD as confirmed by ACS, hx of MI, stable or unstable angina, coronary or other arterial revascularization, stroke,TIA, or peripheral arterial disease presumed to be of atherosclerotic origin. OR C)Primary hyperlipidemia (HLD). HoFH (init): Sub of MR (eg, chart notes, lab values) documenting dx of HoFH as confirmed by one of the following: 1)Gen confirmation of 2 mutations in LDL receptor, ApoB, PCSK9, or LDLRAP1 or ARH, or 2)either untreated LDL greater than 500 or treated LDL greater than 300, AND either xanthoma before 10 yo or evidence of HeFH in both parents. HeFH/ASCVD/Primary HLD (init): One of the following: set A)Both of the following: a)One of the following LDL values while on max tolerated lipid lowering regimen w/in the last 120 days: 1)LDL greater than or equal to 100 mg/dL w/ASCVD, or 2)LDL greater than or equal to 130 mg/dL w/o ASCVD. AND b)One of the following: 1)Pt has been receiving at least 12 wks of one high-intensity (HI) statin therapy (tx) and will continue to receive a HI statin [ie, atorvastatin 40-80 mg, rosuvastatin 20-40 mg] at max tolerated dose.</p>



<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial/Reauth: Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist
<b>Coverage Duration</b>	Initial: 6 months. Reauth: 12 months

Other Criteria	<p>Set A (cont, initial): OR (2) Both of the following: A) Pt unable to tolerate HI statin as evidenced by intolerable and persistent (ie, more than 2 wks) myalgia (muscle symptoms w/o CK elevations) or myositis (muscle symptoms w/ CK elevations less than 10 times ULN) AND B) One of the following: a) Pt has been receiving at least 12 wks of one moderate-intensity (MI) or low-intensity (LI) statin tx and will continue to receive a MI or LI statin [ie, atorvastatin 10-20 mg, rosuvastatin 5-10 mg, simvastatin 10-40 mg, pravastatin 10-80 mg, lovastatin 20-40 mg, Lescol XL (fluvastatin XL) 80 mg, fluvastatin 20-40 mg, or Livalo (pitavastatin) 1-4 mg] at max tolerated dose, OR b) Pt is unable to tolerate MI or LI statin as evidenced by intolerable and persistent (ie, more than 2 weeks) myalgia (muscle symptoms w/o CK elevations) or myositis (muscle symptoms w/ CK elevations less than 10 times ULN), OR (3) Pt has a labeled contraindication to all statins, OR (4) Pt has experienced rhabdomyolysis or muscle symptoms w/ statin treatment w/ CK elevations greater than 10 times ULN on one statin tx. OR set B) Both of the following: a) One of the following LDL values while on max tolerated lipid lowering tx w/in the last 120 days: (1) LDL b/t 70 and 99 mg/dL w/ ASCVD. (2) LDL b/t 100 and 129 mg/dL w/o ASCVD. AND b) Both of the following: (1) One of the following: i) Pt has been receiving at least 12 wks of one max-tolerated statin tx and will continue to receive a statin at max tolerated dose, ii) pt is unable to tolerate statin tx as evidenced by intolerable and persistent (ie, more than 2 wks) myalgia (muscle symptoms w/o CK elevations) or myositis (muscle symptoms w/ CK elevations less than 10 times ULN, iii) Patient has a labeled contraindication to all statins, or iv) Pt has experienced rhabdomyolysis or muscle symptoms w/ statin tx w/ CK elevations greater than 10 times ULN on one statin tx and (2) Pt has been receiving at least 12 weeks of ezetimibe (Zetia) tx as adjunct to max tolerated statin tx OR Pt has a hx of contraindication or intolerance to ezetimibe. HoFH (init): One of the following: 1)Pt is receiving other lipid-lowering tx (eg statin, ezetimibe) or 2)Pt has a documented inability to take other lipid-lowering tx (eg statin, ezetimibe). HeFH/ASCVD/Primary HLD (reauth): Pt continues to receive other lipid-lowering tx (eg statins, ezetimibe) at max tolerated dose (unless pt has documented inability to take these medications). HoFH (reauth): One of the following: 1)Pt continues to receive other lipid-lowering tx (eg statin, ezetimibe) or 2)Pt has a documented inability to take other lipid-lowering tx (eg statin, ezetimibe). HeFH/ASCVD/Primary HLD/HoFH (reauth): Sub of MR (eg, lab values) documenting LDL reduction while on Repatha tx.</p>
----------------	--

## RETEVMO (S)

### Products Affected

- Retevmo

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Lung Cancer: Diagnosis of metastatic non-small cell lung cancer (NSCLC). Disease has presence of RET gene fusion-positive tumor(s). Medullary Thyroid Cancer (MTC): Diagnosis of medullary thyroid cancer (MTC). Disease is advanced or metastatic. Disease has presence of RET gene mutation tumor(s). Disease requires treatment with systemic therapy. Thyroid Cancer: Diagnosis of thyroid cancer. Disease is advanced or metastatic. Disease has presence of RET gene fusion-positive tumor(s). Disease requires treatment with systemic therapy. Patient is radioactive iodine-refractory or radioactive iodine therapy is not appropriate.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Lung Cancer, MTC: Prescribed by or in consultation with an oncologist. Thyroid Cancer: Prescribed by or in consultation with an endocrinologist or an oncologist.
<b>Coverage Duration</b>	Lung Cancer, MTC, Thyroid Cancer: 12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## REVATIO (S)

### Products Affected

- Sildenafil Citrate TABS 20MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
Age Restrictions	N/A
Prescriber Restrictions	PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
Coverage Duration	PAH: Initial: 6 months. Reauth: 12 months.
Other Criteria	PAH (Reauth): Documentation of positive clinical response to therapy.

## REVCovi (S)

---

### Products Affected

- Rvcovi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of adenosine deaminase deficiency (ADA) with severe combined immunodeficiency (SCID).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

## REVLIMID (S)

### Products Affected

- Lenalidomide

- Revlimid

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple myeloma (MM): Diagnosis of MM. Either used as 1) combination therapy with dexamethasone, or 2) maintenance therapy following autologous hematopoietic stem cell transplantation (auto-HSCT). Myelodysplastic syndromes (MDS): Diagnosis of transfusion-dependent anemia due to low- or intermediate-1-risk MDS associated with a deletion 5q. Mantle cell lymphoma (MCL): Diagnosis of MCL. Disease has relapsed after, is refractory to, or progressed after at least one prior therapy (eg, bortezomib, bendamustine, cladribine, rituximab). Follicular Lymphoma (FL): Diagnosis of FL that has been previously treated. Used in combination with a rituximab product. Marginal Zone Lymphoma (MZL): Diagnosis of MZL that has been previously treated. Used in combination with a rituximab product.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist/hematologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## REZUROCK (S)

### Products Affected

- Rezurock

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic graft versus host disease (cGVHD) (initial): Diagnosis of cGVHD. Trial and failure of two or more lines of systemic therapy (e.g., corticosteroids, mycophenolate, etc.).
Age Restrictions	N/A
Prescriber Restrictions	cGVHD (initial): Prescribed by or in consultation with one of the following: hematologist, oncologist, or physician experienced in the management of transplant patients.
Coverage Duration	cGVHD (initial, reauth): 12 months
Other Criteria	cGVHD (reauth): Patient does not show evidence of progressive disease while on therapy.

## RILUTEK (S)

---

### Products Affected

- Riluzole

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Amyotrophic lateral sclerosis (ALS): Diagnosis of amyotrophic lateral sclerosis (ALS).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	ALS: 12 months
Other Criteria	N/A



# RINVOQ (S)

## Products Affected

- Rinvoq

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Rheumatoid arthritis (RA) (initial - 15 mg): Diagnosis of moderately to severely active RA. One of the following: Trial and failure, contraindication, or intolerance (TF/C/I) to one disease modifying antirheumatic drug (DMARD) [eg, methotrexate (Rheumatrex/Trexall), Arava (leflunomide), Azulfidine (sulfasalazine)], OR for continuation of prior therapy. Psoriatic arthritis (PsA) (initial - 15 mg): Diagnosis of active PsA. Ankylosing spondylitis (AS) (initial - 15 mg): Diagnosis of active AS. RA, PsA, AS (initial - 15 mg): Patient has had an inadequate response or intolerance to one or more TNF inhibitors (eg, Enbrel, Humira). Not used in combination with a potent immunosuppressant (e.g., azathioprine, cyclosporine). Atopic dermatitis (AD) (initial - 15 mg and 30 mg): Diagnosis of moderate to severe AD. TF/C/I to at least one of the following: a) Medium or higher potency topical corticosteroid, b) Pimecrolimus cream, c) Tacrolimus ointment, or d) Eucrisa (crisaborole) ointment. One of the following: 1) Trial and failure of a minimum 12-week supply of at least one systemic drug product for the treatment of AD (examples include, but are not limited to, Adbry [tralokinumab-ldrm], Dupixent [dupilumab], etc.), OR 2) Patient has a contraindication, intolerance, or treatment is inadvisable with both of the following FDA-approved AD therapies: Adbry (tralokinumab-ldrm) and Dupixent (dupilumab). Not used in combination with biologic immunomodulators (eg, Dupixent, Adbry) or other immunosuppressants (eg, azathioprine, cyclosporine). Ulcerative colitis (UC) (initial): Diagnosis of moderately to severely active UC. TF/C/I to one of the following conventional therapies: 6-mercaptopurine, aminosalicylate (eg, mesalamine, olsalazine, sulfasalazine), azathioprine, or corticosteroids (eg, prednisone). Patient has had an inadequate response or intolerance to one or more TNF inhibitors (eg, Humira). Not used in combination with a potent immunosuppressant (eg, azathioprine, cyclosporine).</p>

<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	RA, AS (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. AD (initial): Prescribed by or in consultation with a dermatologist or allergist/immunologist. UC (initial): Prescribed by or in consultation with a gastroenterologist.
<b>Coverage Duration</b>	RA, PsA, AS, UC, AD (initial, reauth): 12 months.
<b>Other Criteria</b>	RA, PsA, AS, UC (reauth): Documentation of positive clinical response to therapy. Not used in combination with a potent immunosuppressant (e.g., azathioprine, cyclosporine). AD (reauth): Documentation of a positive clinical response to therapy (eg, reduction in body surface area involvement, reduction in pruritus severity). Not used in combination with biologic immunomodulators (eg, Dupixent, Adbry) or other immunosuppressants (eg, azathioprine, cyclosporine).

## ROZLYTREK (S)

### Products Affected

- Rozlytrek

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic non-small cell lung cancer (NSCLC). Patient has ROS1 rearrangement positive tumor(s). Solid Tumors: Patient has solid tumors with a neurotrophic tyrosine receptor kinase (NTRK) gene fusion (e.g., ETV6-NTRK3, TPM3-NTRK1, TPR-NTRK1, etc.). Disease is without a known acquired resistance mutation (e.g., TRKA G595R, TRKA G667C or TRKC G623R substitutions). Disease is one of the following: metastatic or unresectable (including cases where surgical resection is likely to result in severe morbidity). One of the following: disease has progressed following previous treatment (e.g., surgery, radiation therapy, or systemic therapy) or disease has no satisfactory alternative treatments.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## RUBRACA (S)

### Products Affected

- Rubraca

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Ovarian cancer: Diagnosis of epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. One of the following: 1) Both of the following: a) Presence of deleterious BRCA mutation as detected by a U.S. Food and Drug Administration (FDA)-approved diagnostic test (e.g., FoundationFocus CDxBRCA Assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) and b) Trial and failure, contraindication, or intolerance to two or more chemotherapies (e.g., cisplatin, carboplatin), OR 2) Both of the following: a) Disease is recurrent and b) Used for maintenance treatment in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin). Prostate cancer: Diagnosis of metastatic castration-resistant prostate cancer. Presence of deleterious BRCA mutation as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient has received previous treatment with both of the following: 1) Androgen receptor-directed therapy [e.g., Erleada (apalutamide), Xtandi (enzalutamide), Zytiga (abiraterone)], AND 2) A taxane-based chemotherapy [e.g., docetaxel, Jevtana (cabazitaxel)].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Ovarian cancer: Prescribed by or in consultation with an oncologist. Prostate cancer: Prescribed by or in consultation with an oncologist or urologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy

# RUXIENCE (S)

## Products Affected

- Ruxience

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-Hodgkin's Lymphoma (NHL): One of the following: 1) Diagnosis of follicular, CD20-positive, B-cell non-Hodgkin's lymphoma. Used as first-line treatment in combination with chemotherapy, 2) Diagnosis of follicular, CD20-positive, B-cell non-Hodgkin's lymphoma. Patient achieved a complete or partial response to a rituximab product in combination with chemotherapy. Used as monotherapy for maintenance therapy, 3) Diagnosis of low-grade, CD20-positive, B-cell non-Hodgkin's lymphoma. One of the following: a) Patient has stable disease following first-line treatment with CVP (cyclophosphamide, vincristine, prednisolone/ prednisone) chemotherapy or, b) Patient achieved a partial or complete response following first-line treatment with CVP (cyclophosphamide, vincristine, prednisolone/ prednisone) chemotherapy, 4) Diagnosis of relapsed or refractory, low grade or follicular CD20-positive, B-cell non-Hodgkin's lymphoma OR, 5) Diagnosis of diffuse large B-cell, CD20-positive, non-Hodgkin's lymphoma. Used as first-line treatment in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) or other anthracycline-based chemotherapy regimens. Chronic Lymphocytic Leukemia (CLL): Diagnosis of chronic lymphocytic leukemia. Used in combination with fludarabine and cyclophosphamide. Wegener's Granulomatosis (WG) and Microscopic Polyangiitis (MPA): Diagnosis of WG or MPA. Patient is concurrently on glucocorticoids (eg, prednisone) OR contraindication or intolerance to glucocorticoids (eg, prednisone).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	NHL, CLL: Prescribed by or in consultation with a hematologist/oncologist. WG, MPA: Prescribed by or in consultation with a nephrologist, pulmonologist, or rheumatologist.
<b>Coverage Duration</b>	NHL, CLL: 12 months. WG, MPA: 3 months.

<b>Other Criteria</b>	Approve for continuation of prior therapy.
-----------------------	--

# RYDAPT (S)

## Products Affected

- Rydapt

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute Myeloid Leukemia (AML): Newly diagnosed acute myeloid leukemia (AML), FMS-like tyrosine kinase 3 (FLT3) mutation-positive as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., LeukoStrat CDx FLT3 Mutation Assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA), used in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation. Aggressive Systemic Mastocytosis (ASM), Systemic Mastocytosis with Associated Hematological Neoplasm (SM-AHN), Mast Cell Leukemia (MCL): Diagnosis of one of the following: aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated hematological neoplasm (SM-AHN), or mast cell leukemia (MCL).
Age Restrictions	N/A
Prescriber Restrictions	All uses: Prescribed by or in consultation with a hematologist or oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## SABRIL (S)

### Products Affected

- Vigabatrin
- Vigadrone

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Complex Partial Seizures (CPS): For use as adjunctive therapy. Failure, contraindication, or intolerance to two formulary anticonvulsants [eg, Lamictal (lamotrigine), Depakene (valproic acid), Dilantin (phenytoin)]. Infantile Spasms (IS): Diagnosis of infantile spasms.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.



# SANDOSTATIN (S)

## Products Affected

- Octreotide Acetate INJ 1000MCG/ML, 100MCG/ML, 200MCG/ML, 500MCG/ML, 50MCG/ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acromegaly (initial): Diagnosis of acromegaly. One of the following: A) Inadequate response to surgical resection and/or pituitary irradiation OR B) Patient is not a candidate for surgical resection or pituitary irradiation. Trial and failure, contraindication or intolerance to a dopamine agonist (e.g., bromocriptine or cabergoline) at maximally tolerated doses. Carcinoid tumor (initial): Diagnosis of metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea or flushing episodes. Vasoactive intestinal peptide tumor (initial): Diagnosis of vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	All uses (initial, reauth): 12 months
Other Criteria	Acromegaly (reauth): Documentation of positive clinical response to therapy (e.g., reduction or normalization of IGF-1/GH level for same age and sex, reduction in tumor size). Carcinoid tumor (reauth): Patient has improvement in number of diarrhea or flushing episodes. Vasoactive intestinal peptide tumor (reauth): Patient has improvement in number of diarrhea episodes.

## SARCLISA (S)

---

### Products Affected

- Sarclisa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of multiple myeloma. Patient has received at least two prior treatment regimens which included lenalidomide and a proteasome inhibitor (e.g., bortezomib, carfilzomib). Used in combination with pomalidomide and dexamethasone.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist/hematologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## SCSEMBLIX (S)

### Products Affected

- Scemblix

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of chronic myelogenous/myeloid leukemia (CML). Disease is Philadelphia chromosome-positive (Ph+). Disease is in chronic phase. One of the following: 1) Patient has been previously treated with two or more alternative tyrosine kinase inhibitors (TKI) [e.g., Bosulif (bosutinib), imatinib, Sprycel (dasatinib), Tassigna (nilotinib), Iclusig (ponatinib)], OR 2) Disease is T315I mutation positive.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or hematologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## SCIG (S)

### Products Affected

- Cuvitru

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	All uses (initial, reauth): Contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication).
<b>Required Medical Information</b>	Initial: Immune globulin will be administered at the minimum effective dose and appropriate frequency for the prescribed diagnosis. Medication is being used subcutaneously. Diagnosis of chronic inflammatory demyelinating polyneuropathy (CIDP) OR one of the following FDA-approved or literature supported diagnoses: 1) Common variable immunodeficiency (CVID), OR 2) Congenital agammaglobulinemia (X-linked or autosomal recessive), OR 3) Severe combined immunodeficiencies (SCID), OR 4) Wiskott-Aldrich syndrome, OR 5) Other primary immunodeficiency with an immunologic evaluation including IgG levels below the normal laboratory value for the patient's age at the time of diagnosis and the patient lacks an adequate response to protein and polysaccharide antigens (i.e., tetanus toxoid or diphtheria toxoid and pneumovax or HiB vaccine).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses (initial, reauth): Prescribed by or in consultation with a physician who has specialized expertise in managing patients on SCIG therapy (e.g., immunologist, hematologist, neurologist).
<b>Coverage Duration</b>	Initial, reauth: 12 months
<b>Other Criteria</b>	Subject to Part B vs. Part D review. All uses (reauth): Patient has experienced an objective improvement on immune globulin therapy and the immune globulin will be administered at the minimum effective dose (by decreasing the dose, increasing the frequency, or implementing both strategies) for maintenance therapy.

## SECUADO (S)

### Products Affected

- Secuado

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of schizophrenia. Both of the following: 1) Trial and failure of Saphris (asenapine) and 2) Trial and failure, contraindication, or intolerance to one of the following generic formulary atypical antipsychotic agents: aripiprazole, olanzapine, paliperidone, quetiapine, risperidone, or ziprasidone.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## SIGNIFOR (S)

### Products Affected

- Signifor

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cushing's disease (initial): Diagnosis of Cushing's disease. One of the following: a) Pituitary surgery has not been curative for the patient or b) Patient is not a candidate for pituitary surgery.
Age Restrictions	N/A
Prescriber Restrictions	Cushing's disease (initial): Prescribed by or in consultation with an endocrinologist.
Coverage Duration	Cushing's disease (initial, reauth): 12 months
Other Criteria	Cushing's disease (reauth): Documentation of positive clinical response to therapy (e.g., a clinically meaningful reduction in 24-hour urinary free cortisol levels, improvement in signs or symptoms of the disease).

## SKYRIZI (S)

### Products Affected

- Skyrizi
- Skyrizi Pen

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. Psoriatic arthritis (PsA) (Initial): Diagnosis of active PsA.
Age Restrictions	N/A
Prescriber Restrictions	Plaque psoriasis (initial): Prescribed by or in consultation with a dermatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist.
Coverage Duration	Plaque psoriasis, PsA (Initial, reauth): 12 months
Other Criteria	Plaque psoriasis (Reauth): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline. PsA (Reauth): Documentation of positive clinical response to therapy.

# SOMATULINE DEPOT (S)

## Products Affected

- Somatuline Depot

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acromegaly: Diagnosis of acromegaly. One of the following: A) Inadequate response to one of the following: surgery or radiotherapy, OR B) Not a candidate for one of the following: surgery or radiotherapy. Gastroenteropancreatic neuroendocrine tumors (GEP-NETs) (120mg/0.5mL strength only): Diagnosis of GEP-NETs. Disease is one of the following: (a) unresectable, locally advanced or (b) metastatic. Carcinoid syndrome (120mg/0.5mL strength only): Diagnosis of carcinoid syndrome. Used to reduce the frequency of short-acting somatostatin analog rescue therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	All uses: 12 months
Other Criteria	All Indications: Approve for continuation of prior therapy.



## SOMAVERT (S)

### Products Affected

- Somavert

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acromegaly (initial): Diagnosis of acromegaly AND Failure to surgery and/or radiation therapy and/or other medical therapies (such as dopamine agonists [e.g., bromocriptine, cabergoline]) unless patient is not a candidate for these treatment options AND trial and failure or intolerance to generic octreotide (a somatostatin analogue)
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	Initial and reauth: 12 months
Other Criteria	Acromegaly (reauth): Patient has experienced a positive clinical response to therapy (biochemical control, decrease or normalization of IGF-1 levels).

# SPORANOX (S)

## Products Affected

- Itraconazole CAPS
- Itraconazole SOLN

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	One of the following: 1) patient has a systemic fungal infection (e.g., aspergillosis, histoplasmosis, blastomycosis), OR 2) all of the following: a) patient has a diagnosis of onychomycosis confirmed by one of the following (CAPSULE ONLY): i) positive potassium hydroxide (KOH) preparation, OR ii) fungal culture, OR iii) nail biopsy, AND b) patient has had a trial and failure, contraindication, or intolerance to oral terbinafine, OR 3) both of the following (ORAL SOLUTION ONLY): a) patient has a diagnosis of candidiasis (esophageal or oropharyngeal), AND b) one of the following: i) candidiasis is refractory or resistant to treatment with fluconazole OR ii) trial and failure, contraindication, or intolerance to fluconazole.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Systemic fungal infxn:6mo.Candidiasis:1mo.Fingernail onycho:5wks.Toenail onycho, other:3mo.
Other Criteria	N/A

## SPRAVATO (S)

### Products Affected

- Spravato 56mg Dose
- Spravato 84mg Dose

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	One of the following: A) Both of the following: 1) Diagnosis of major depressive disorder (treatment-resistant) and 2) Patient has not experienced a clinical meaningful improvement after treatment with at least two antidepressants from different classes for an adequate duration (at least 4 weeks each) in the current depressive episode OR B) Both of the following: 1) Diagnosis of major depressive disorder and 2) Patient has both of the following: a) depressive symptoms and b) acute suicidal ideation or behavior. Used in combination with an oral antidepressant (e.g., duloxetine, escitalopram, sertraline).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a psychiatrist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## SPRYCEL (S)

### Products Affected

- Sprycel

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Philadelphia chromosome positive (Ph+)/BCR ABL chronic myelogenous leukemia (CML): Diagnosis of Ph+/BCR ABL CML. Ph+/BCR ABL acute lymphoblastic leukemia (ALL): Diagnosis of Ph+/BCR ABL ALL.
Age Restrictions	N/A
Prescriber Restrictions	All Uses: Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	All Uses: 12 months
Other Criteria	All Uses: Approve for continuation of prior therapy.

## STELARA (IV) (S)

### Products Affected

- Stelara INJ 130MG/26ML

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Crohn's Disease (CD): Diagnosis of moderately to severely active CD. Both of the following: trial and failure, contraindication, or intolerance to Humira (adalimumab), and trial and failure, contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, Imuran (azathioprine), methotrexate, corticosteroid (eg, prednisone, methylprednisolone). Ulcerative Colitis (UC): Diagnosis of moderately to severely active UC. Trial and failure, contraindication, or intolerance to both Humira (adalimumab) and Xeljanz/Xeljanz XR (tofacitinib).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist.
<b>Coverage Duration</b>	One time
<b>Other Criteria</b>	Stelara is to be administered as an intravenous induction dose. Stelara induction dosing is in accordance with the United States Food and Drug Administration approved labeled dosing for Crohn's Disease/ulcerative colitis: 260 mg for patients weighing 55 kg or less, 390 mg for patients weighing more than 55 kg to 85 kg, or 520 mg for patients weighing more than 85 kg.

# STELARA (S)

## Products Affected

- Stelara INJ 45MG/0.5ML, 90MG/ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Plaque psoriasis (Initial - 45mg/0.5mL): Diagnosis of moderate to severe plaque psoriasis. Plaque psoriasis (Initial - 90mg/1mL): Diagnosis of moderate to severe plaque psoriasis. Patient's weight is greater than 100 kg (220 lbs). Plaque psoriasis (Initial): Trial and failure, contraindication (eg, safety concerns, not indicated for patient's age/weight), or intolerance to two of the following: Enbrel (etanercept), Humira (adalimumab), Skyrizi (risankizumab), Cosentyx (secukinumab), OR for continuation of prior therapy. Psoriatic arthritis (PsA) (Initial - 45mg/0.5mL): Diagnosis of active PsA. PsA (Initial - 90mg/1mL): Diagnosis of active PsA. Patient's weight is greater than 100 kg (220 lbs). Diagnosis of co-existent moderate to severe psoriasis. PsA (Initial): Trial and failure, contraindication, or intolerance (TF/C/I) to two of the following, or attestation demonstrating a trial may be inappropriate: Enbrel (etanercept), Humira (adalimumab), Skyrizi (risankizumab), Rinvoq (upadacitinib), Xeljanz/XR (tofacitinib/ER), OR for continuation of prior therapy. Crohn's disease (CD) (Initial): Diagnosis of moderately to severely active Crohn's disease. Both of the following: TF/C/I to Humira (adalimumab) and TF/C/I to treatment with at least one immunomodulator or corticosteroid [e.g., Purinethol (6-mercaptopurine), Imuran (azathioprine), Sandimmune (cyclosporine A), Prograf (tacrolimus), MTX (methotrexate), prednisone, methylprednisolone], OR for continuation of prior therapy. Ulcerative colitis (UC) (Initial): Diagnosis of moderately to severely active UC. TF/C/I to two of the following, or attestation demonstrating a trial may be inappropriate: Humira (adalimumab), Rinvoq (upadacitinib), or Xeljanz/Xeljanz XR (tofacitinib), OR for continuation of prior therapy.</p>
Age Restrictions	N/A

<b>Prescriber Restrictions</b>	Plaque psoriasis (initial): Prescribed by or in consultation with a dermatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. CD and UC (initial): Prescribed by or in consultation with a gastroenterologist.
<b>Coverage Duration</b>	All uses (Initial, reauth): 12 months
<b>Other Criteria</b>	Reauth (PsA, CD, UC): Documentation of positive clinical response to therapy. Reauth (plaque psoriasis): Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, OR improvement in symptoms (eg, pruritus, inflammation) from baseline.

# STIVARGA (S)

## Products Affected

- Stivarga

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Metastatic colorectal cancer (mCRC): All of the following: 1) diagnosis of mCRC, AND 2) trial and failure, contraindication or intolerance to fluoropyrimidine-, oxaliplatin- and irinotecan-based chemotherapy (e.g., FOLFOX, FOLFIRI, FOLFOXIRI), AND 3) trial and failure, contraindication or intolerance to an anti-VEGF therapy (e.g., bevacizumab), AND 4) one of the following: a) RAS mutation, OR b) both of the following: RAS wild-type (RAS mutation negative tumor) and trial and failure, contraindication or intolerance to an anti-EGFR therapy [e.g., Vectibix (panitumumab), Erbitux (cetuximab)]. Gastrointestinal stromal tumor (GIST): All of the following: 1) diagnosis of locally advanced, unresectable or metastatic GIST, AND 2) trial and failure, contraindication or intolerance to both of the following: a) Gleevec (imatinib mesylate), AND b) Sutent (sunitinib malate). Hepatocellular Carcinoma (HCC): Diagnosis of HCC. Trial and failure or intolerance to Nexavar (sorafenib tosylate).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	mCRC, GIST: Prescribed by or in consultation with an oncologist. HCC: Prescribed by or in consultation with an oncologist, hepatologist, or gastroenterologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.



## STRENSIQ (S)

### Products Affected

- Strensiq

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Hypophosphatasia: Diagnosis of perinatal/infantile or juvenile-onset hypophosphatasia AND for patients requesting the 80 mg/0.8 mL vial only: Patient's weight is greater than or equal to 40 kg.
Age Restrictions	N/A
Prescriber Restrictions	Hypophosphatasia: Prescribed by or in consultation with a specialist experienced in the treatment of inborn errors of metabolism or endocrinologist
Coverage Duration	Hypophosphatasia: 12 months
Other Criteria	N/A

## SUPPRELIN LA (S)

### Products Affected

- Supprelin La

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Central Precocious Puberty (CPP) (initial): Diagnosis of CPP (idiopathic or neurogenic). Early onset of secondary sexual characteristics in females less than age 8 or males less than age 9. Advanced bone age of at least one year compared with chronologic age. One of the following: a) patient has undergone gonadotropin-releasing hormone agonist (GnRHa) testing AND Peak luteinizing hormone (LH) level above pre-pubertal range, or b) patient has a random LH level in the pubertal range.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CPP (initial, reauth): Prescribed by or in consultation with a pediatric endocrinologist.
<b>Coverage Duration</b>	CPP (init, reauth): 12 months
<b>Other Criteria</b>	CPP (reauthorization): LH levels have been suppressed to pre-pubertal levels.

## SUTENT (S)

### Products Affected

- Sunitinib Malate
- Sutent

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Renal cell carcinoma: Diagnosis of advanced or metastatic renal cell carcinoma. Gastrointestinal stromal tumor (GIST): Diagnosis of GIST after disease progression on, or contraindication or intolerance to Gleevec (imatinib). Pancreatic neuroendocrine tumors: Diagnosis of progressive, well-differentiated pancreatic neuroendocrine tumor that is unresectable locally advanced or metastatic disease. Adjuvant treatment of renal cell carcinoma: Diagnosis of renal cell carcinoma (RCC). Used as adjuvant therapy. Patient is at high risk of recurrent RCC following nephrectomy.
Age Restrictions	N/A
Prescriber Restrictions	All uses: Prescribed by or in consultation with an oncologist
Coverage Duration	All uses: 12 months
Other Criteria	All Indications: Approve for continuation of prior therapy.

## SYLATRON (S)

---

### Products Affected

- Sylatron

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Melanoma: Diagnosis of melanoma with microscopic or gross nodal involvement AND The prescribed medication will be used as adjuvant therapy within 84 days of definitive surgical resection, including complete lymphadenectomy.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# SYNAGIS (S)

## Products Affected

- Synagis INJ 100MG/ML, 50MG/0.5ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Patient will use palivizumab for immunoprophylaxis of respiratory syncytial virus (RSV) during the peak months of infection in the patient's geographic region AND Patient meets one of the following criteria: 1) Infants born at 28 weeks, six days gestation or earlier and who are younger than 12 months of age at the start of the RSV season OR 2) Diagnosis of chronic lung disease of prematurity, born before 32 weeks, 0 days gestation, received greater than 21% oxygen for at least the first 28 days after birth, and one of the following: a) 12 months of age or younger at the start of the RSV season OR b) greater than 12 months of age to 24 months of age at the start of the RSV season and received medical support (i.e., chronic corticosteroid therapy, diuretic therapy, or supplemental oxygen) within 6 months before the start of the second RSV season. OR 3) Patient is 12 months of age or younger at the start of the RSV season and has one of the following: a) acyanotic heart failure that will require a cardiac surgical procedure and the patient is receiving medication to control congestive heart failure, OR b) moderate to severe pulmonary hypertension OR c) cyanotic heart defect. OR 4) patient is younger than 24 months of age and will or has undergone a cardiac transplantation during the RSV season. OR 5) Patient is 12 months of age or younger at the start of the RSV season with a congenital abnormality or neuromuscular disorder and has an impaired ability to clear secretions from the upper airway due to an ineffective cough. OR 6) Patient is younger than 24 months of age with a lymphocyte count below the normal range for patient's age and has received or will receive a solid organ transplant, hematopoietic stem cell transplant recipient, or chemotherapy during the RSV season.</p>
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a pediatric specialist (i.e., pulmonologist, neonatologist, neurologist, cardiologist, pediatric intensivist, or infectious disease specialist).

<b>Coverage Duration</b>	5 months (5 doses) during RSV season.
<b>Other Criteria</b>	N/A

## SYNRIBO (S)

### Products Affected

- Synribo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic myelogenous leukemia (CML): Diagnosis of CML in the chronic or accelerated phase AND Patient has tried and has had resistance, relapse, inadequate response, intolerance or is contraindicated to TWO tyrosine kinase inhibitors (i.e., Gleevec [imatinib], Sprycel, Tasigna, and Bosulif, Iclusig).
Age Restrictions	N/A
Prescriber Restrictions	CML: Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## SYPRINE (S)

### Products Affected

- Clovique
- Trientine Hydrochloride

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial: Diagnosis of Wilson's disease (i.e., hepatolenticular degeneration). Trial and failure, contraindication, or intolerance to a penicillamine product (e.g., Depen, Cuprimine)
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Reauth: Documentation of a positive clinical response to therapy



## TABRECTA (S)

### Products Affected

- Tabrecta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of non-small cell lung cancer (NSCLC). Disease is one of the following: recurrent, advanced, metastatic. Presence of mesenchymal-epithelial transition (MET) exon 14 skipping positive tumors as detected with an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# TAFAMIDIS (S)

## Products Affected

- Vyndaqel

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM) (initial): Diagnosis of transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM). One of the following: 1) Patient has a transthyretin (TTR) mutation (e.g., V122I), 2) Cardiac or noncardiac tissue biopsy demonstrating histologic confirmation of TTR amyloid deposits, OR 3) All of the following: i) echocardiogram or cardiac magnetic resonance imaging suggestive of amyloidosis, ii) scintigraphy scan suggestive of cardiac TTR amyloidosis, and iii) absence of light-chain amyloidosis. One of the following: 1) History of heart failure (HF), with at least one prior hospitalization for HF, OR 2) Presence of clinical signs and symptoms of HF (e.g., dyspnea, edema). Patient has New York Heart Association (NYHA) Functional Class I, II, or III heart failure.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	ATTR-CM (initial, reauth): Prescribed by or in consultation with a cardiologist
<b>Coverage Duration</b>	ATTR-CM (initial, reauth): 12 months
<b>Other Criteria</b>	ATTR-CM (reauth): Documentation of positive clinical response to therapy. Patient continues to have New York Heart Association (NYHA) Functional Class I, II, or III heart failure.

# TAFINLAR (S)

## Products Affected

- Tafinlar

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Melanoma: Diagnosis of unresectable or metastatic melanoma AND cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) OR both of the following: cancer is BRAF V600E or V600K mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) and medication is used in combination with Mekinist (trametinib). Adjuvant Treatment for Melanoma: Diagnosis of melanoma. Cancer is BRAF V600E or V600K mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Involvement of lymph nodes following complete resection. Used as adjunctive therapy. Medication is used in combination with Mekinist (trametinib). Non-small Cell Lung Cancer (NSCLC): Diagnosis of metastatic non-small cell lung cancer AND cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) AND medication is used in combination with Mekinist (trametinib). Anaplastic Thyroid Cancer (ATC): Diagnosis of locally advanced or metastatic anaplastic thyroid cancer. Cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Cancer may not be treated with standard locoregional treatment options. Medication is used in combination with Mekinist (trametinib) .</p>
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.

<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

# TAGRISSO (S)

## Products Affected

- Tagrisso

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): One of the following: A) All of the following: Diagnosis of metastatic NSCLC. One of the following: 1) Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA), OR 2) Both of the following: a) Patient has known active EGFR T790M mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) and b) Patient has experienced disease progression on or after one of the following EGFR Tyrosine Kinase Inhibitors (TKIs): Gilotrif (afatinib), Iressa (gefitinib), Tarceva (erlotinib), or Vizimpro (dacomitinib). OR B) All of the following: Diagnosis of NSCLC. Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Both of the following: 1) Patient is receiving as adjuvant therapy, and 2) Patient has had a complete surgical resection of the primary NSCLC tumor.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## TALZENNA (S)

### Products Affected

- Talzenna

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: Diagnosis of breast cancer. Disease is one of the following: a) locally advanced or b) metastatic. Presence of a deleterious or suspected deleterious germline BRCA-mutation (gBRCAm) as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Disease is human epidermal growth factor receptor 2 (HER2)-negative.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# TARCEVA (S)

## Products Affected

- Erlotinib Hydrochloride

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of locally advanced or metastatic (Stage III or IV) NSCLC AND Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Pancreatic Cancer: Diagnosis of locally advanced, unresectable, or metastatic pancreatic cancer AND erlotinib will be used in combination with gemcitabine.
Age Restrictions	N/A
Prescriber Restrictions	All uses: Prescribed by or in consultation with an oncologist
Coverage Duration	All uses: 12 months
Other Criteria	All Indications: Approve for continuation of prior therapy.

# TARGRETIN (S)

## Products Affected

- Bexarotene

- Targretin GEL

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cutaneous T-Cell Lymphoma (CTCL): Diagnosis of CTCL. Trial and failure, contraindication, or intolerance to at least one prior therapy (including skin-directed therapies [eg, corticosteroids {ie, clobetasol, diflorasone, halobetasol, augmented betamethasone dipropionate}] or systemic therapies [eg, interferons]).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.



# TARPEYO (S)

## Products Affected

- Tarpeyo

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of primary immunoglobulin A nephropathy (IgAN). Patient is at risk of rapid disease progression [e.g., generally a urine protein-to-creatinine ratio (UPCR) greater than or equal to 1.5 g/g, or by other criteria such as clinical risk scoring using the International IgAN Prediction Tool]. Used to reduce proteinuria. Estimated glomerular filtration rate (eGFR) greater than or equal to 35 mL/min/1.73 m <sup>2</sup> . One of the following: 1) Patient has been on a minimum 90-day trial of a maximally tolerated dose and will continue to receive therapy with one of the following: a) an angiotensin-converting enzyme (ACE) inhibitor (e.g., benazepril, lisinopril), or b) an angiotensin II receptor blocker (ARB) (e.g., losartan, valsartan), OR 2) Patient has a contraindication or intolerance to both ACE inhibitors and ARBs. Trial and failure, contraindication, or intolerance to another glucocorticoid (e.g., methylprednisolone, prednisone).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a nephrologist.
<b>Coverage Duration</b>	9 months.
<b>Other Criteria</b>	N/A

## TASIGNA (S)

---

### Products Affected

- Tasigna

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic myelogenous leukemia (CML): Diagnosis of Ph+/BCR ABL CML
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# TAVALISSE (S)

## Products Affected

- Tavalisse

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic Idiopathic Thrombocytopenic Purpura (ITP) (initial): Diagnosis of chronic immune ITP or relapsed/refractory ITP. Baseline platelet count is less than 30,000/mcL. Trial and failure, contraindication, or intolerance to at least one of the following: corticosteroids (e.g., prednisone, methylprednisolone), immunoglobulins [e.g., Gammagard, immune globulin (human)], splenectomy, thrombopoietin receptor agonists (e.g., Nplate, Promacta), or Rituxan (rituximab). Patient's degree of thrombocytopenia and clinical condition increase the risk of bleeding.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	ITP (initial): Prescribed by or in consultation with a hematologist/oncologist.
<b>Coverage Duration</b>	ITP (initial, reauth): 12 months
<b>Other Criteria</b>	ITP (reauth): Documentation of positive clinical response to therapy as evidenced by an increase in platelet count to a level sufficient to avoid clinically important bleeding.

# TAVNEOS (S)

## Products Affected

- Tavneos

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Initial: Diagnosis of one of the following types of severe active anti-neutrophil cytoplasmic autoantibody (ANCA)-associated vasculitis: a) Granulomatosis with polyangiitis (GPA) OR b) Microscopic polyangiitis (MPA). Diagnosis is confirmed by one of the following: a) ANCA test positive for proteinase 3 (PR3) antigen, b) ANCA test positive for myeloperoxidase (MPO) antigen, OR c) Tissue biopsy. Patient is receiving concurrent immunosuppressant therapy with one of the following: a) cyclophosphamide OR b) rituximab. One of the following: a) Patient is concurrently on glucocorticoids (e.g., prednisone) OR b) History of contraindication or intolerance to glucocorticoids (e.g., prednisone).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial, Reauth: Prescribed by or in consultation with a nephrologist, pulmonologist, or rheumatologist
<b>Coverage Duration</b>	Initial, Reauth: 12 months
<b>Other Criteria</b>	Reauth: Patient does not show evidence of progressive disease while on therapy. Patient is receiving concurrent immunosuppressant therapy (e.g., azathioprine, cyclophosphamide, methotrexate, rituximab).

## TAZVERIK (S)

### Products Affected

- Tazverik

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Epithelioid sarcoma: Diagnosis of epithelioid sarcoma. Disease is one of the following: metastatic or locally advanced. Patient is not eligible for complete resection. Follicular lymphoma: Diagnosis of follicular lymphoma. Disease is one of the following: relapsed or refractory.
Age Restrictions	N/A
Prescriber Restrictions	Epithelioid sarcoma: Prescribed by or in consultation with an oncologist. Follicular lymphoma: Prescribed by or in consultation with an oncologist or hematologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# TECFIDERA (S)

## Products Affected

- Dimethyl Fumarate CPDR
- Dimethyl Fumarate Starterpack MISC 0

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple Sclerosis (MS) (initial): Diagnosis of a relapsing form of MS (eg, clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions).
Age Restrictions	N/A
Prescriber Restrictions	MS (initial, reauth): Prescribed by or in consultation with a neurologist
Coverage Duration	MS (initial, reauth): 12 months
Other Criteria	MS (reauth): Documentation of positive clinical response to therapy (e.g., stability in radiologic disease activity, clinical relapses, disease progression).

# TEGSEDI (S)

## Products Affected

- Tegsedi

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Hereditary transthyretin-mediated amyloidosis (hATTR amyloidosis) (initial): Diagnosis of hATTR amyloidosis with polyneuropathy. Patient has a transthyretin (TTR) mutation (e.g., V30M). One of the following: 1) Patient has a baseline polyneuropathy disability (PND) score less than or equal to IIIb, 2) Patient has baseline familial amyloidotic polyneuropathy (FAP) stage of 1 or 2, OR 3) Patient has a baseline neuropathy impairment score (NIS) between 10 and 130. Presence of clinical signs and symptoms of the disease (e.g., peripheral/autonomic neuropathy).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	hATTR amyloidosis (initial): Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	hATTR amyloidosis (initial, reauth): 12 months
<b>Other Criteria</b>	hATTR amyloidosis (reauth): Patient has demonstrated a benefit from therapy (e.g., improved neurologic impairment, slowing of disease progression, quality of life assessment). One of the following: 1) Patient continues to have a PND score less than or equal to IIIb, 2) Patient continues to have a FAP stage of 1 or 2, OR 3) Patient continues to have a NIS between 10 and 130.

## TEPMETKO (S)

---

### Products Affected

- Tepmetko

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Disease is metastatic. Presence of mesenchymal-epithelial transition (MET) exon 14 skipping alterations.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.



# TERIPARATIDE (S)

## Products Affected

- Forteo INJ 600MCG/2.4ML
- Teriparatide

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Postmenopausal osteoporosis or osteopenia or men with primary or hypogonadal osteoporosis or osteopenia (initial): Diagnosis of one of the following: a) postmenopausal osteoporosis or osteopenia or b) primary or hypogonadal osteoporosis or osteopenia. One of the following: Set I) Both of the following: A) Bone mineral density (BMD) T-score of -2.5 or lower in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) trial and failure, contraindication, or intolerance (TF/C/I) to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]), or Set II) Both of the following: A) BMD T-score between -1.0 and -2.5 in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) both of the following: i) TF/C/I to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]) and ii) One of the following FRAX 10-year probabilities: a) Major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) Hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions. Glucocorticoid-Induced Osteoporosis: See Other Criteria section.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	All uses (initial): 24 months. All uses (reauth): 12 months.

<b>Other Criteria</b>	<p>Glucocorticoid-Induced Osteoporosis (initial): Diagnosis of glucocorticoid-induced osteoporosis. History of prednisone or its equivalent at a dose greater than or equal to 5mg/day for greater than or equal to 3 months. One of the following: 1) BMD T-score less than or equal to -2.5 based on BMD measurements from lumbar spine, femoral neck, total hip, or radius (one-third radius site), or 2) One of the following FRAX 10-year probabilities: a) Major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) Hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions, or 3) History of one of the following fractures resulting from minimal trauma: vertebral compression fx, fx of the hip, fx of the distal radius, fx of the pelvis, or fx of the proximal humerus. TF/C/I to one bisphosphonate (e.g., alendronate). All uses (initial, reauth): One of the following: 1) Treatment duration of parathyroid hormones [e.g., teriparatide, Tymlos (abaloparatide)] has not exceeded a total of 24 months during the patient's lifetime, or 2) Patient remains at or has returned to having a high risk for fracture despite a total of 24 months of use of parathyroid hormones [e.g., teriparatide, Tymlos (abaloparatide)].</p>
-----------------------	--

# TESTOSTERONE (S)

## Products Affected

- Androderm PT24 2MG/24HR, 4MG/24HR
- Testosterone GEL 25MG/2.5GM, 50MG/5GM
- Testosterone Cypionate INJ 100MG/ML, 200MG/ML
- Testosterone Pump

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Hypogonadism (HG) (Initial): Diagnosis (dx) of HG AND male patient at birth AND one of the following: 1) Two pre-treatment serum total testosterone (T) levels less than 300 ng/dL (10.4 nmol/L) or less than the reference range for the lab, OR 2) Both of the following: a) Has a condition that may cause altered sex-hormone binding globulin (SHBG) (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and b) one pre-treatment calculated free or bioavailable T level less than 5 ng/dL (0.17 nmol/L) or less than reference range for the lab, OR 3) History of bilateral orchiectomy, panhypopituitarism, or a genetic disorder known to cause HG (eg, congenital anorchia, Klinefelter's syndrome), OR 4) Both of the following: a) Patient is continuing testosterone therapy, and b) One of the following: i) Follow-up total serum T level or calculated free or bioavailable T level drawn within the past 12 months is within or below the normal limits of the reporting lab, or ii) follow-up total serum T level or calculated free or bioavailable T level drawn within the past 12 months is outside of upper limits of normal for the reporting lab and the dose is adjusted. Gender Dysphoria (GD)/Gender Incongruence (off-label): Dx of GD/Gender Incongruence. Patient is a transgender male (female-to-male).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	HG(init): (New to T tx:6 mo. New to plan and cont T tx:12 mo), (reauth): 12 mo. GD: 12 mo.

<b>Other Criteria</b>	<p>HG (Reauth): 1) Follow-up total serum T level within or below the normal limits of the reporting lab, or 2) Follow-up total serum T level outside of upper limits of normal for the reporting lab and the dose is adjusted, OR 3) Has a condition that may cause altered SHBG (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and one of the following: Follow-up calculated free or bioavailable T level within or below the normal limits of the reporting lab, or follow-up calculated free or bioavailable T level outside of upper limits of normal for the reporting lab and the dose is adjusted.</p>
-----------------------	---

# TESTOSTERONE ENANTHATE (S)

## Products Affected

- Testosterone Enanthate INJ

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Hypogonadism (HG) (Initial): Diagnosis (dx) of HG AND male patient at birth AND one of the following: 1) Two pre-treatment serum total testosterone (T) levels less than 300 ng/dL (10.4 nmol/L) or less than the reference range for the lab, OR 2) Both of the following: a) Has a condition that may cause altered sex-hormone binding globulin (SHBG) (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and b) one pre-treatment calculated free or bioavailable T level less than 5 ng/dL (0.17 nmol/L) or less than reference range for the lab, OR 3) History of bilateral orchiectomy, panhypopituitarism, or a genetic disorder known to cause HG (eg, congenital anorchia, Klinefelter's syndrome), OR 4) Both of the following: a) Patient is continuing testosterone therapy, and b) One of the following: i) Follow-up total serum T level or calculated free or bioavailable T level drawn within the past 12 months is within or below the normal limits of the reporting lab, or ii) follow-up total serum T level or calculated free or bioavailable T level drawn within the past 12 months is outside of upper limits of normal for the reporting lab and the dose is adjusted. Delayed puberty (DP): Dx of DP AND male patient at birth. Breast cancer (BC): Dx of inoperable BC AND used for palliative treatment AND female patient at birth. Gender Dysphoria (GD)/Gender Incongruence (off-label): Dx of GD/Gender Incongruence. Patient is a transgender male (female-to-male).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	HG(init): (New to T tx:6 mo. Cont T tx:12 mo), (reauth): 12 mo. BC, GD: 12 mo. DP: 6 mo.

<b>Other Criteria</b>	<p>HG (Reauth): 1) Follow-up total serum T level within or below the normal limits of the reporting lab, or 2) Follow-up total serum T level outside of upper limits of normal for the reporting lab and the dose is adjusted, OR 3) Has a condition that may cause altered SHBG (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and one of the following: Follow-up calculated free or bioavailable T level within or below the normal limits of the reporting lab, or follow-up calculated free or bioavailable T level outside of upper limits of normal for the reporting lab and the dose is adjusted.</p>
-----------------------	---

## THALOMID (S)

---

### Products Affected

- Thalomid

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple myeloma (MM): Diagnosis of MM. Used in combination with dexamethasone, unless the patient has an intolerance to steroids. Erythema nodosum leprosum (ENL): Diagnosis of moderate to severe ENL with cutaneous manifestations. Thalomid is not used as monotherapy if moderate to severe neuritis is present.
Age Restrictions	N/A
Prescriber Restrictions	MM: Prescribed by or in consultation with an oncologist/hematologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# TIBSOVO (S)

## Products Affected

- Tibsovo

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Relapsed or refractory Acute Myeloid Leukemia (AML): Diagnosis of AML. Disease is relapsed or refractory. Newly-Diagnosed AML: Diagnosis of newly-diagnosed AML. One of the following: 1) patient is greater than or equal to 75 years old OR 2) patient has comorbidities that preclude use of intensive induction chemotherapy. Locally Advanced or Metastatic Cholangiocarcinoma: Diagnosis of cholangiocarcinoma. Disease is locally advanced or metastatic. Patient has been previously treated. All indications: Patient has an isocitrate dehydrogenase-1 (IDH1) mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., Abbott RealTime IDH1 assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Relapsed or refractory AML, Newly-Diagnosed AML: Prescribed by or in consultation with a hematologist/oncologist. Locally Advanced or Metastatic Cholangiocarcinoma: Prescribed by or in consultation with a hepatologist, gastroenterologist, or oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.



## TOPICAL RETINOID (S)

---

### Products Affected

- Tretinoin CREA 0.025%, 0.05%

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acne vulgaris: Diagnosis of acne vulgaris (i.e., acne).
Age Restrictions	PA applies to members 26 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

# TRAZIMERA (S)

## Products Affected

- Trazimera

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast cancer: Diagnosis of human epidermal growth factor receptor 2 (HER2)-overexpressing breast cancer. One of the following treatment regimens: a) As adjuvant treatment, b) metastatic disease and one of the following: 1) used in combination with a taxane (eg, docetaxel, paclitaxel), or 2) used as a single agent in a patient who has received one or more chemotherapy regimens for metastatic disease, or c) used in combination with Perjeta (pertuzumab). Gastric Cancer: Diagnosis of HER2-overexpressing gastric or gastroesophageal junction adenocarcinoma (locally advanced, recurrent, or metastatic). Used in combination with one of the following treatment regimens: a) Platinol (cisplatin) and Adrucil (5-fluorouracil), or b) Platinol (cisplatin) and Xeloda (capecitabine).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## TRELSTAR (S)

### Products Affected

- Trelstar Mixject INJ 11.25MG, 22.5MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prostate Cancer: Diagnosis of advanced or metastatic prostate cancer. Trial and failure, contraindication, or intolerance to any brand Lupron formulation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## TRIKAFTA (S)

### Products Affected

- Trikafta TBPK 50MG; 0; 25MG

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cystic Fibrosis (CF) (initial): Diagnosis of CF. Patient has at least one of the following mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene as detected by an FDA-cleared cystic fibrosis mutation test or a test performed at a Clinical Laboratory Improvement Amendments (CLIA)-approved facility: F508del mutation OR a mutation in the CFTR gene that is responsive based on in vitro data.
<b>Age Restrictions</b>	CF (initial): 6 years of age or older.
<b>Prescriber Restrictions</b>	CF (initial): Prescribed by or in consultation with a pulmonologist or specialist affiliated with a CF care center.
<b>Coverage Duration</b>	CF (initial, reauth): 12 months
<b>Other Criteria</b>	CF (reauth): Documentation of positive clinical response to therapy (e.g., improvement in lung function [percent predicted forced expiratory volume in one second {PPFEV1}] or decreased number of pulmonary exacerbations).

# TRIPTODUR (S)

## Products Affected

- Triptodur

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Central Precocious Puberty (CPP) (initial): Diagnosis of CPP (idiopathic or neurogenic). Early onset of secondary sexual characteristics in females less than age 8 or males less than age 9. Advanced bone age of at least one year compared with chronologic age. One of the following: a) patient has undergone gonadotropin-releasing hormone agonist (GnRHa) testing AND Peak luteinizing hormone (LH) level above pre-pubertal range, or b) patient has a random LH level in the pubertal range. Trial and failure or intolerance to Lupron Depot-Ped.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CPP (initial, reauth): Prescribed by or in consultation with a pediatric endocrinologist.
<b>Coverage Duration</b>	CPP (Initial, reauth): 12 months
<b>Other Criteria</b>	CPP (reauthorization): LH levels have been suppressed to pre-pubertal levels.

## TRODELVY (S)

### Products Affected

- Trodelvy

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Triple Negative Breast Cancer (TNBC): Diagnosis of TNBC. Disease is metastatic. Patient has received at least two prior therapies for metastatic disease (e.g., carboplatin, cisplatin, gemcitabine, paclitaxel, docetaxel, capecitabine, etc.).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## TRUSELTIQ (S)

### Products Affected

- Truseltiq

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of cholangiocarcinoma. Disease is one of the following: a) unresectable locally advanced or b) metastatic. Disease has presence of a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient has been previously treated.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hepatologist, gastroenterologist, or oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## TUKYSA (S)

### Products Affected

- Tukysa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: Diagnosis of breast cancer. Disease is one of the following: a) advanced unresectable or b) metastatic. Disease is human epidermal growth factor receptor 2 (HER2)-positive. Used in combination with trastuzumab and capecitabine. Patient has received one or more prior anti-HER2 based regimens (e.g., trastuzumab, pertuzumab, ado-trastuzumab emtansine).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.



## TURALIO (S)

---

### Products Affected

- Turalio

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Tenosynovial Giant Cell Tumor (TGCT): Diagnosis of TGCT. Patient is symptomatic. Patient is not a candidate for surgery due to worsening functional limitation or severe morbidity with surgical removal.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## TYKERB (S)

### Products Affected

- Lapatinib Ditosylate

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast Cancer: Diagnosis of human epidermal growth factor receptor 2 (HER2)-positive metastatic or recurrent breast cancer. Used in combination with one of the following: Trastuzumab, Xeloda (capecitabine), or aromatase inhibitors [eg, Aromasin (exemestane), Femara (letrozole), Arimidex (anastrozole)].
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# TYMLOS (S)

## Products Affected

- Tymlos

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of postmenopausal osteoporosis or osteopenia. One of the following: Set I) Both of the following: A) Bone mineral density (BMD) T-score of -2.5 or lower in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) trial and failure, contraindication, or intolerance (TF/C/I) to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]), or Set II) Both of the following: A) BMD T-score between -1.0 and -2.5 in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) both of the following: i) TF/C/I to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]) and ii) one of the following FRAX (Fracture Risk Assessment Tool) 10-year probabilities: a) major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions. Treatment duration of parathyroid hormones (e.g., teriparatide, Tymlos [abaloparatide]) has not exceeded a total of 24 months during the patient's lifetime.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	24 months (max 24 months of therapy per lifetime)
Other Criteria	N/A

# TYSABRI (S)

## Products Affected

- Tysabri

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Multiple Sclerosis (MS) (initial): Diagnosis of a relapsing form of MS (eg, clinically isolated syndrome, relapsing-remitting disease, secondary progressive disease, including active disease with new brain lesions). One of the following: 1) Trial and failure, contraindication, or intolerance (TF/C/I) to one of the following disease-modifying therapies for MS: A) Aubagio (teriflunomide), B) Lemtrada (alemtuzumab), C) Mavenclad (cladribine), D) Plegridy (peginterferon beta-1a), E) Any one of the inteferon beta-1a injections (eg, Avonex), F) Any one of the interferon beta-1b injections (eg, Betaseron, Extavia), G) Any one of the glatiramer acetate injections (eg, Copaxone, Glatopa, generic glatiramer acetate), H) Any one of the oral fumarates (eg, brand Tecfidera, generic dimethyl fumarate), I) Any one of the Sphingosine 1-Phosphate (S1P) receptor modulators (eg, Gilenya, Mayzent, Zeposia), J) Any one of the B-cell targeted therapies (eg, Ocrevus, Kesimpta), 2) Patient is not a candidate for any of the drugs listed as prerequisites due to the severity of their MS, or 3) for continuation of prior therapy. MS (init, reauth): Not used in combination with another disease-modifying therapy for MS. Crohn's Disease (CD) (initial): Diagnosis of moderately to severely active CD with evidence of inflammation (eg, elevated C-reactive protein [CRP], elevated erythrocyte sedimentation rate, presence of fecal leukocytes). TF/C/I to one of the following conventional therapies: corticosteroids (eg, prednisone, methylprednisolone), 6-mercaptopurine (6MP [Purinethol], azathioprine (Imuran), methotrexate (Rheumatrex, Trexall), aminosalicylates (eg, sulfasalazine, mesalamine, olsalazine). TF/C/I to a TNF-inhibitor (eg, Humira [adalimumab], infliximab). CD (initial and reauth): Not used in combination with an immunosuppressant (eg, 6-MP, azathioprine, cyclosporine, or methotrexate). Not used in combination with a TNF-inhibitor (eg, Enbrel [etanercept], Humira [adalimumab], or infliximab).</p>
Age Restrictions	N/A

<b>Prescriber Restrictions</b>	MS (init, reauth): Prescribed by or in consultation with a neurologist. CD (initial): Prescribed by or in consultation with a gastroenterologist.
<b>Coverage Duration</b>	MS (init, reauth): 12mo. CD (Init): 3 mo. CD (Reauth): 6 mo if not on steroids. Otherwise, 3 mo.
<b>Other Criteria</b>	MS (reauth): Documentation of positive clinical response to therapy (e.g., stability in radiologic disease activity, clinical relapses, disease progression). CD (reauth): Documentation of positive clinical response (eg, improved disease activity index) to therapy.

## UBRELVY (S)

### Products Affected

- Ubrelvy

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial: Diagnosis of migraine with or without aura. Will be used for the acute treatment of migraine. Will not be used for preventive treatment of migraine. Patient has fewer than 15 headache days per month. Trial and failure or intolerance to one triptan (e.g., eletriptan, rizatriptan, sumatriptan) or a contraindication to all triptans. Medication will not be used in combination with another oral CGRP inhibitor.
Age Restrictions	Initial: 18 years of age or older.
Prescriber Restrictions	Initial, Reauth: Prescribed by or in consultation with a neurologist, headache specialist, or pain specialist.
Coverage Duration	Initial: 3 months. Reauth: 12 months.
Other Criteria	Reauth: Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea). Will not be used for preventive treatment of migraine. Medication will not be used in combination with another oral CGRP inhibitor.

# UDENYCA (S)

## Products Affected

- Udenyca

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Febrile neutropenia (FN) prophylaxis: Patient will be receiving prophylaxis for FN due to one of the following: 1) Patient is receiving National Cancer Institute's Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer, 2) patient is receiving a dose-dense chemotherapy regimen for which the incidence of FN is unknown, 3) patient is receiving chemotherapy regimen(s) associated with greater than 20% incidence of FN, 4) both of the following: a) patient is receiving chemotherapy regimen(s) associated with 10-20% incidence of FN, AND b) patient has one or more risk factors associated with chemotherapy-induced infection, FN, or neutropenia, OR 5) Both of the following: a) patient is receiving myelosuppressive anticancer drugs associated with neutropenia, AND b) patient has a history of FN or dose-limiting event during a previous course of chemotherapy (secondary prophylaxis). Treatment of FN (off-label): Patient has received or is receiving myelosuppressive anticancer drugs associated with neutropenia. Diagnosis of FN. Patient is at high risk for infection-associated complications. Acute radiation syndrome (ARS) (off-label): Patient was/will be acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of ARS).</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with a hematologist/oncologist
<b>Coverage Duration</b>	ARS: 1 mo. FN (prophylaxis, treatment): 3 mo or duration of tx.
<b>Other Criteria</b>	N/A

# UKONIQ (S)

## Products Affected

- Ukoniq

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Marginal zone lymphoma (MZL): Diagnosis of MZL. Disease is one of the following: relapsed or refractory. Patient has received at least one prior anti-CD20-based regimen (e.g., bendamustine + rituximab, bendamustine + obinutuzumab, etc.). Follicular lymphoma (FL): Diagnosis of FL. Disease is one of the following: relapsed or refractory. Patient has received at least three prior lines of systemic therapy (e.g., bendamustine + rituximab, bendamustine + obinutuzumab, etc.).
Age Restrictions	N/A
Prescriber Restrictions	MZL/FL: Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.



## VALCHLOR (S)

### Products Affected

- Valchlor

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Mycosis fungoides-type cutaneous T-cell lymphoma (MF-CTCL) (initial): All of the following: 1) diagnosis of Stage IA MF-CTCL, OR diagnosis of Stage IB MF-CTCL, AND 2) patient has received at least one prior skin-directed therapy [e.g., topical corticosteroids, bexarotene topical gel (Targretin topical gel), etc.].
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## VARIZIG (S)

### Products Affected

- Varizig INJ 125UNIT/1.2ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Presence of contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication).
Required Medical Information	Immune globulin is being used intramuscularly. The immune globulin is being used for passive immunization or post exposure-prophylaxis of varicella. Patient is considered a high risk individual (i.e., immune compromised, pregnant woman, newborn of mother with varicella, premature infant, and infant less than 1 year old).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months (approve one dose only)
Other Criteria	N/A

## VASCEPA (S)

### Products Affected

- Icosapent Ethyl

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Severe Hypertriglyceridemia (init): Diagnosis (dx) of hypertriglyceridemia and patient has a pre-treatment triglyceride (TG) level greater than or equal to 500 mg/dL. Prevention of CV Events (init): Dx of hypertriglyceridemia and patient has a pre-treatment TG level of 150 to 499 mg/dL. One of the following: 1) Patient has established cardiovascular disease (CVD) (e.g., coronary artery disease, cerebrovascular or carotid disease, peripheral artery disease, etc.) OR 2) Both of the following: a) Dx of diabetes mellitus AND b) Patient has two or more risk factors for developing CVD. Medication will be used as an adjunct to maximally tolerated statin therapy unless there is a contraindication or intolerance to statin therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial/Reauth: 12 months
<b>Other Criteria</b>	Severe Hypertriglyceridemia (reauth): Documentation of positive clinical response to therapy. Prevention of CV Events (Reauth): Documentation of positive clinical response to therapy. Medication continues to be used as an adjunct to maximally tolerated statin therapy unless there is a contraindication or intolerance to statin therapy.

## VENCLEXTA (S)

### Products Affected

- Venclexta

- Venclexta Starting Pack

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL): Diagnosis of CLL or SLL. Acute Myeloid Leukemia (AML): Diagnosis of newly diagnosed AML. Used in combination with azacitidine, or decitabine, or low-dose cytarabine. One of the following: 1) age 75 years or older OR 2) comorbidities that preclude use of intensive induction chemotherapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## VENTAVIS (S)

### Products Affected

- Ventavis

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
Age Restrictions	N/A
Prescriber Restrictions	PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist.
Coverage Duration	PAH (Initial): 6 months. (Reauth): 12 months
Other Criteria	Subject to Part B vs D review. PAH (Reauth): Documentation of positive clinical response to therapy.

# VERZENIO (S)

## Products Affected

- Verzenio

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Advanced or Metastatic Breast Cancer: Diagnosis of advanced or metastatic breast cancer. Disease is hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative. One of the following: a) used in combination with an aromatase inhibitor (e.g., Arimidex [anastrozole], Aromasin [exemestane], Femara [letrozole]) and patient is a postmenopausal woman, OR b) used in combination with Faslodex (fulvestrant) and disease has progressed following endocrine therapy, OR c) used as monotherapy and disease has progressed following endocrine therapy and patient has already received at least one prior chemotherapy regimen. Early Breast Cancer: Diagnosis of early breast cancer at high risk of recurrence. Disease is hormone receptor (HR)-positive. Disease is human epidermal growth factor receptor 2 (HER2)-negative. Disease is node-positive. Used as adjunctive therapy. Used in combination with one of the following endocrine therapies: 1) tamoxifen or 2) aromatase inhibitor (e.g., anastrozole, letrozole, exemestane). Patient has a Ki-67 score of greater than or equal to 20% as determined by an FDA approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## VIMIZIM (S)

### Products Affected

- Vimizim

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Mucopolysaccharidosis (initial): Diagnosis of Mucopolysaccharidosis type IVA (MPS IVA, Morquio A syndrome) confirmed by both of the following: a) documented clinical signs and symptoms of the disease (e.g., kyphoscoliosis, genu valgum, pectus carinatum, gait disturbance, growth deficiency, etc.) and b) documented reduced fibroblast or leukocyte GALNS enzyme activity or molecular genetic testing of GALNS.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial, reauth: 12 months
Other Criteria	Reauth: Documentation of positive clinical response to therapy.

## VITRAKVI (S)

### Products Affected

- Vitrakvi

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Presence of solid tumors (e.g., salivary gland, soft tissue sarcoma, infantile fibrosarcoma, thyroid cancer, lung, melanoma, colon, etc.).</p> <p>Disease is positive for neurotrophic receptor tyrosine kinase (NTRK) gene fusion (e.g. ETV6-NTRK3, TPM3-NTRK1, LMNA-NTRK1, etc.).</p> <p>Disease is without a known acquired resistance mutation [e.g., TRKA G595R substitution, TRKA G667C substitution, or other recurrent kinase domain (solvent front and xDFG) mutations]. Disease is one of the following: metastatic or unresectable (including cases where surgical resection is likely to result in severe morbidity). One of the following:</p> <p>Disease has progressed on previous treatment (e.g., surgery, radiotherapy, or systemic therapy) OR Disease has no satisfactory alternative treatments.</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.



## VIZIMPRO (S)

---

### Products Affected

- Vizimpro

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Disease is metastatic. Disease is positive for one of the following epidermal growth factor receptor (EGFR) mutations: exon 19 deletion or exon 21 L858R substitution.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# VORICONAZOLE INJECTION (S)

## Products Affected

- Voriconazole INJ

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Invasive aspergillosis: Diagnosis of invasive aspergillosis (IA). Candidemia: Diagnosis of candidemia. One of the following: (1) patient is non-neutropenic or (2) infection is located in skin, abdomen, kidney, bladder wall, or wounds. Esophageal Candidiasis: Diagnosis of esophageal candidiasis. Mycosis: Diagnosis of fungal infection caused by <i>Scedosporium apiospermum</i> (asexual form of <i>Pseudallescheria boydii</i> ) or <i>Fusarium</i> spp. including <i>Fusarium solani</i> . For fusariosis: Patient is intolerant of, or refractory to, other therapy (e.g., liposomal amphotericin B, amphotericin B lipid complex).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 weeks
Other Criteria	N/A

## VOSEVI (S)

### Products Affected

- Vosevi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guideline. All patients: Diagnosis of chronic hepatitis C, patient is without decompensated liver disease (defined as Child-Pugh Class B or C), and not used in combination with another HCV direct acting antiviral agent [e.g., Harvoni, Zepatier].
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine.
Coverage Duration	12 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline.
Other Criteria	N/A

## VOTRIENT (S)

### Products Affected

- Votrient

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Renal cell carcinoma (RCC): Diagnosis of advanced/metastatic RCC. Soft tissue sarcoma: Diagnosis of advanced soft tissue sarcoma and patient received at least one prior chemotherapy (e.g., ifosfamide, doxorubicin, cisplatin, dacarbazine, docetaxel, oxaliplatin, etc.).
Age Restrictions	N/A
Prescriber Restrictions	All uses: Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## WELIREG (S)

### Products Affected

- Welireg

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of von Hippel-Lindau (VHL) disease. Patient requires therapy for one of the following: a) renal cell carcinoma (RCC), b) central nervous system (CNS) hemangioblastoma, or c) pancreatic neuroendocrine tumor (pNET). Patient does not require immediate surgery.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## XALKORI (S)

### Products Affected

- Xalkori

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Non-small cell lung cancer (NSCLC): Diagnosis of advanced or metastatic NSCLC AND One of the following: A) Patient has an anaplastic lymphoma kinase (ALK)-positive tumor as detected with a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) or B) Patient has MET amplification- or ROS1 rearrangement-positive tumor as detected with an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Anaplastic Large Cell Lymphoma (ALCL): Diagnosis of systemic ALCL. Disease is relapsed or refractory. Patient has an anaplastic lymphoma kinase (ALK)-positive tumor as detected with a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	NSCLC: Prescribed by or in consultation with an oncologist. ALCL: Prescribed by or in consultation with an oncologist or hematologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## XCOPRI (S)

---

### Products Affected

- Xcopri

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of partial onset seizures.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# XELJANZ (S)

## Products Affected

- Xeljanz
- Xeljanz Xr

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Xeljanz tab/Xeljanz XR tab: Rheumatoid arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Trial and failure, contraindication, or intolerance (TF/C/I) to one disease modifying antirheumatic drug (DMARD) [eg, methotrexate (Rheumatrex/Trexall), Arava (leflunomide), Azulfidine (sulfasalazine)]. Psoriatic arthritis (PsA) (Initial): Diagnosis of active PsA. RA, PsA (Initial): Patient has had an inadequate response or intolerance to one or more TNF inhibitors (eg, Enbrel, Humira). Ankylosing spondylitis (AS) (Initial): Diagnosis of active AS. One of the following: TF/C/I to Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior tofacitinib therapy. Ulcerative colitis (UC) (Initial): Diagnosis of moderately to severely active UC. Trial and failure, contraindication or intolerance to one of the following conventional therapies: 6-mercaptopurine (Purinethol), aminosalicylate [e.g., mesalamine (Asacol, Pentasa, Rowasa), olsalazine (Dipentum), sulfasalazine (Azulfidine, Sulfazine)], azathioprine (Imuran), or corticosteroids (e.g., prednisone, methylprednisolone). Patient has had an inadequate response or intolerance to one or more TNF inhibitors (eg, Humira). All indications: Patient is not receiving tofacitinib in combination with a potent immunosuppressant (eg, azathioprine, cyclosporine).
Age Restrictions	N/A
Prescriber Restrictions	RA, PJI, AS (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. UC (initial): Prescribed by or in consultation with a gastroenterologist.
Coverage Duration	RA/PJI/PSA/AS (initial, reauth): 12 months. UC (init): 4 mo. UC (reauth): 12 mo.



<b>Other Criteria</b>	<p>Xeljanz: Polyarticular course juvenile idiopathic arthritis (PJIA) (Initial): Diagnosis of active polyarticular course juvenile idiopathic arthritis. One of the following: TF/C/I to Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR patient has a documented needle-phobia to the degree that the patient has previously refused any injectable therapy or medical procedure (refer to DSM-V-TR 300.29/F40.2 for specific phobia diagnostic criteria), OR for continuation of prior therapy. All Indications (Reauth): Documentation of positive clinical response to therapy. Patient is not receiving tofacitinib in combination with a potent immunosuppressant (eg, azathioprine, cyclosporine).</p>
-----------------------	--

## XENAZINE (S)

### Products Affected

- Tetrabenazine

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chorea associated with Huntington's Disease (HD) (Initial): Diagnosis of chorea in patients with Huntington's disease. Tardive dyskinesia (Initial): Diagnosis of tardive dyskinesia. One of the following: 1) Patient has persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering, or discontinuation of the offending medication, OR 2) Patient is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication. Tourette's syndrome (Initial): Patient has tics associated with Tourette's syndrome. Trial and failure, contraindication, or intolerance to Haldol (haloperidol).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	HD (Initial): Prescribed by or in consultation with a neurologist. Tardive dyskinesia, Tourette's syndrome (Initial): Prescribed by or in consultation with neurologist or psychiatrist.
<b>Coverage Duration</b>	All uses: (initial) 3 months. (Reauth) 12 months.
<b>Other Criteria</b>	All indications (Reauth): Documentation of clinical response and benefit from therapy.

## XERMELO (S)

### Products Affected

- Xermelo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Carcinoid syndrome diarrhea (Initial): Diagnosis of carcinoid syndrome diarrhea AND diarrhea is inadequately controlled by a stable dose of somatostatin analog (SSA) therapy (e.g., octreotide [Sandostatin, Sandostatin LAR], lanreotide [Somatuline Depot]) for at least 3 months AND used in combination with SSA therapy.
Age Restrictions	N/A
Prescriber Restrictions	Initial: Prescribed by or in consultation with an oncologist, endocrinologist, or gastroenterologist
Coverage Duration	Initial: 6 months. Reauth: 12 months
Other Criteria	Carcinoid syndrome diarrhea (Reauthorization): Documentation of a positive clinical response to therapy AND drug will continue to be used in combination with SSA therapy.

## XGEVA (S)

### Products Affected

- Xgeva

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Multiple Myeloma (MM)/Bone metastasis from solid tumors (BMST): One of the following: 1) Both of the following: a) Diagnosis of multiple myeloma and b) Trial and failure, intolerance, or contraindication to one bisphosphonate (eg, zoledronic acid), OR 2) Both of the following: a) Diagnosis of solid tumors (eg, breast cancer, kidney cancer, lung cancer, prostate cancer, thyroid cancer) and b) Documented evidence of one or more metastatic bone lesions. Giant cell tumor of bone (GCTB): Both of the following: 1) Diagnosis of giant cell tumor of bone AND 2) One of the following: a) tumor is unresectable, OR b) surgical resection is likely to result in severe morbidity. Hypercalcemia of malignancy (HCM): Both of the following: 1) Diagnosis of hypercalcemia of malignancy, AND 2) Trial and failure, contraindication, or intolerance to one intravenous bisphosphonate (eg, pamidronate, Zometa (zoledronic acid)).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	GCTB, HCM: Prescribed by or in consultation with an oncologist
<b>Coverage Duration</b>	MM/BMST, GCTB: 12 mo. HCM: 2 mo.
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## XIFAXAN (S)

### Products Affected

- Xifaxan

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Travelers' diarrhea (TD) (200 mg strength only): Diagnosis of travelers' diarrhea, AND one of the following: a) Trial and failure, contraindication, or intolerance to one of the following: Cipro (ciprofloxacin), Levaquin (levofloxacin), ofloxacin, Zithromax (azithromycin) OR b) resistance to all of the following: Cipro (ciprofloxacin), Levaquin (levofloxacin), ofloxacin, Zithromax (azithromycin). Prophylaxis of hepatic encephalopathy (HE) recurrence (550mg strength only): Used for the prophylaxis of hepatic encephalopathy recurrence, AND trial and failure, contraindication or intolerance to lactulose. Treatment of HE: Used for the treatment of HE. Trial and failure, contraindication, or intolerance to lactulose. Irritable bowel syndrome with diarrhea (IBS-D) (550mg strength only) (initial): Diagnosis of IBS-D, AND trial and failure, contraindication or intolerance to an antidiarrheal agent [eg, loperamide].
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	TD: 14 days. HE (prophylaxis, treatment): 12 months. IBS-D (initial, reauth): 2 weeks.
<b>Other Criteria</b>	IBS-D (reauth): Patient experiences IBS-D symptom recurrence.

# XOLAIR (S)

## Products Affected

- Xolair

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Asthma (init): Diagnosis of moderate to severe persistent allergic asthma. Positive skin test or in vitro reactivity to a perennial aeroallergen. Pretreatment serum immunoglobulin (Ig)E level between 30 to 700 IU/mL for patients 12 years of age and older OR 30 to 1300 IU/mL for patients 6 years to less than 12 years of age. Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) High-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and ii) additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium), OR b) One maximally-dosed combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone/vilanterol)].</p> <p>Chronic Idiopathic Urticaria (CIU) (init): Diagnosis of CIU. Persistent symptoms (itching and hives) for at least 4 consecutive weeks despite titrating to an optimal dose with a second generation H1 antihistamine (e.g., cetirizine, fexofenadine), unless there is a contraindication or intolerance to H1 antihistamines. Patient has tried and had an inadequate response or intolerance or contraindication to at least one of the following additional therapies: H2 antagonist (e.g., famotidine, cimetidine), leukotriene receptor antagonist (e.g., montelukast), H1 antihistamine, hydroxyzine, doxepin. Used concurrently with an H1 antihistamine, unless there is a contraindication or intolerance to H1 antihistamines.</p> <p>Nasal polyps (NP) (init): Diagnosis of NP. Unless contraindicated, the patient has had an inadequate response to 2 months of treatment with an intranasal corticosteroid (e.g., fluticasone, mometasone). Used in combination with another agent for nasal polyps.</p>
Age Restrictions	N/A

<b>Prescriber Restrictions</b>	Asthma (init/reauth): Prescribed by or in consultation with an allergist/immunologist, or pulmonologist. CIU (init): Prescribed by or in consultation with an allergist/immunologist, or dermatologist. NP (init/reauth): Prescribed by or in consultation with an allergist/immunologist, otolaryngologist, or pulmonologist.
<b>Coverage Duration</b>	Asthma, init: 6 mo, reauth: 12 mo. CIU, init: 3 mo, reauth: 6 mo. NP, init/reauth: 12 mo.
<b>Other Criteria</b>	Asthma (reauth): Documentation of positive clinical response to therapy (e.g., Reduction in asthma exacerbations, improvement in forced expiratory volume in 1 second (FEV1), decreased use of rescue medications). Patient continues to be treated with an inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) with or without additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium) unless there is a contraindication or intolerance to these medications. CIU (reauth): Patient's disease status has been re-evaluated since the last authorization to confirm the patient's condition warrants continued treatment. Patient has experienced one or both of the following: Reduction in itching severity from baseline or Reduction in the number of hives from baseline. NP (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in nasal polyps score [NPS: 0-8 scale], improvement in nasal congestion/obstruction score [NCS: 0-3 scale]). Used in combination with another agent for nasal polyps.

## XOSPATA (S)

---

### Products Affected

- Xospata

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of acute myeloid leukemia (AML). Disease is relapsed or refractory. Patient has a FMS-like tyrosine kinase (FLT3) mutation as determined by a U.S. Food and Drug Administration (FDA)-approved test (e.g., LeukoStrat CDx FLT3 Mutation Assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.



# XPOVIO (S)

## Products Affected

- Xpovio
- Xpovio 100 Mg Once Weekly
- Xpovio 40 Mg Once Weekly
- Xpovio 40 Mg Twice Weekly
- Xpovio 60 Mg Once Weekly
- Xpovio 60 Mg Twice Weekly
- Xpovio 80 Mg Once Weekly
- Xpovio 80 Mg Twice Weekly

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple Myeloma (MM): Diagnosis of multiple myeloma. Patient has received at least one prior therapy (e.g., lenalidomide, bortezomib, daratumumab, pomalidomide). Used in combination with bortezomib and dexamethasone. Relapsed/Refractory Multiple Myeloma (RRMM): Diagnosis of relapsed or refractory multiple myeloma (RRMM). Patient has received at least four prior therapies (e.g., lenalidomide, bortezomib, daratumumab, pomalidomide). Disease is refractory to all of the following: 1) Two proteasome inhibitors (e.g., bortezomib, carfilzomib), 2) Two immunomodulatory agents (e.g., lenalidomide, thalidomide), and 3) An anti-CD38 monoclonal antibody (e.g. daratumumab). Used in combination with dexamethasone. Diffuse large B-cell lymphoma (DLBCL): Diagnosis of one of the following: 1) Relapsed or refractory DLBCL not otherwise specified OR 2) Relapsed or refractory DLBCL arising from follicular lymphoma. Patient has previously received at least two lines of systemic therapy (e.g., CHOP: cyclophosphamide, doxorubicin, vincristine, and prednisone plus rituximab).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist/hematologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## XTANDI (S)

### Products Affected

- Xtandi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Castration-resistant or castration-recurrent prostate cancer (CRPC): Diagnosis of castration-resistant (chemical or surgical) or recurrent prostate cancer. Metastatic castration-sensitive prostate cancer (M-CSPC): Diagnosis of metastatic castration-sensitive prostate cancer.
Age Restrictions	N/A
Prescriber Restrictions	CRPC, M-CSPC: Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# XYREM (S)

## Products Affected

- Xyrem

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Narcolepsy with cataplexy (Narcolepsy Type 1)(initial): Diagnosis of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible), AND symptoms of cataplexy are present, AND symptoms of excessive daytime sleepiness (eg, irrepressible need to sleep or daytime lapses into sleep) are present. Narcolepsy without cataplexy (Narcolepsy Type 2)(initial): Diagnosis of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible), AND symptoms of cataplexy are absent, AND symptoms of excessive daytime sleepiness (eg, irrepressible need to sleep or daytime lapses into sleep) are present, AND trial and failure, contraindication, or intolerance to one of the following: 1) amphetamine-based stimulant (eg, amphetamine, dextroamphetamine), OR 2) methylphenidate-based stimulant.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses (initial): Prescribed by or in consultation with one of the following: neurologist, psychiatrist, or sleep medicine specialist.
<b>Coverage Duration</b>	All uses (initial): 6 months. All uses (reauth): 12 months
<b>Other Criteria</b>	Narcolepsy Type 1 (reauth): Documentation demonstrating a reduction in the frequency of cataplexy attacks associated with therapy, OR documentation demonstrating a reduction in symptoms of excessive daytime sleepiness associated with therapy. Narcolepsy Type 2 (reauth): Documentation demonstrating a reduction in symptoms of excessive daytime sleepiness associated with therapy.

## ZAVESCA (S)

---

### Products Affected

- Miglustat

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Gaucher disease: Diagnosis of mild to moderate type 1 Gaucher disease. Patient is unable to receive enzyme replacement therapy due to one of the following conditions: allergy or hypersensitivity to enzyme replacement therapy, poor venous access, unavailability of enzyme replacement therapy (e.g., Cerezyme, VPRIV).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Gaucher disease: 12 months
Other Criteria	N/A

# ZEJULA (S)

## Products Affected

- Zejula

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of one of the following: advanced epithelial ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Used for maintenance treatment in patients who are in a complete or partial response to first-line platinum-based chemotherapy (e.g., cisplatin, carboplatin). Recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of one of the following: recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer. Used for maintenance treatment in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin). Treatment of advanced ovarian cancer after three or more chemotherapies: Diagnosis of advanced ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Patient has been treated with three or more prior chemotherapy regimens. Patient's cancer is associated with homologous recombination deficiency (HRD) positive status defined by one of the following: (a) a deleterious or suspected deleterious BRCA mutation or (b) both of the following: (1) genomic instability and (2) cancer has progressed more than 6 months after response to the last platinum-based chemotherapy (e.g., cisplatin, carboplatin).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## ZELBORAF (S)

### Products Affected

- Zelboraf

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Melanoma: Diagnosis of unresectable melanoma or metastatic melanoma. Cancer is BRAFV600 mutant type (MT) as detected by a U.S. Food and Drug Administration (FDA)-approved test (eg, cobas 4600 BRAFV600 Mutation Test) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Erdheim-Chester Disease: Diagnosis of Erdheim-Chester disease AND Disease is BRAFV600 mutant type (MT).
Age Restrictions	N/A
Prescriber Restrictions	Melanoma: Prescribed by or in consultation with an oncologist. Erdheim-Chester Disease: Prescribed by or in consultation with a hematologist/oncologist.
Coverage Duration	12 months
Other Criteria	All indications: Approve for continuation of therapy.

## ZEPZELCA (S)

---

### Products Affected

- Zepzelca

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of metastatic small cell lung cancer (SCLC). Disease has progressed on or after platinum-based chemotherapy (e.g., carboplatin, cisplatin, oxaliplatin).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# ZIRABEV (S)

## Products Affected

- Zirabev

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Non-Small Cell Lung Cancer: Excluded if squamous cell histology.
Required Medical Information	<p>Colorectal Cancer: Diagnosis of metastatic colorectal cancer. One of the following: 1) Both of the following: a) used as first- or second-line treatment and b) used in combination with an intravenous 5-fluorouracil-based chemotherapy, OR 2) All of the following: a) used as second-line treatment, b) used in combination with fluoropyrimidine-irinotecan-based chemotherapy or fluoropyrimidine-oxaliplatin-based chemotherapy, and c) patient has progressed on a first-line bevacizumab-containing regimen.</p> <p>Non-Small Cell Lung Cancer (NSCLC): Diagnosis of NSCLC. Disease is unresectable, locally advanced, recurrent, or metastatic. Used as first-line treatment. Used in combination with paclitaxel and carboplatin.</p> <p>Renal Cell Cancer: Diagnosis of metastatic renal cell cancer. Used in combination with interferon-alpha.</p> <p>Cervical Cancer: Diagnosis of carcinoma of the cervix. Disease is persistent, recurrent, or metastatic. Used in combination with one of the following: a) paclitaxel and cisplatin or b) paclitaxel and topotecan.</p> <p>Glioblastoma: Diagnosis of recurrent glioblastoma.</p> <p>Epithelial Ovarian, Fallopian Tube, or Primary Peritoneal Cancer: Diagnosis of epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. One of the following: 1) All of the following: a) disease is stage 3 or 4, b) patient has been treated with bevacizumab as a single agent, c) treatment is following surgical resection, and d) used in combination with carboplatin and paclitaxel, OR 2) All of the following: a) disease is platinum-resistant recurrent, b) patient has received no more than 2 prior chemotherapy regimens, and c) used in combination with paclitaxel, pegylated liposomal doxorubicin, or topotecan, OR 3) All of the following: a) disease is platinum-sensitive recurrent, b) patient has been treated with bevacizumab as a single agent, and c) used in combination with one of the following: i) carboplatin and paclitaxel or ii) carboplatin and gemcitabine.</p>
Age Restrictions	N/A



<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Hepatocellular Carcinoma (HCC): Diagnosis of hepatocellular carcinoma. Disease is unresectable or metastatic. Used in combination with Tecentriq (atezolizumab). Patient has not received prior systemic therapy. Approve for continuation of prior therapy.

## ZOKINVY (S)

### Products Affected

- Zokinvy

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	One of the following: 1) Diagnosis of Hutchinson-Gilford Progeria Syndrome, OR 2) For treatment of processing-deficient Progeroid Laminopathies with one of the following: i) Heterozygous LMNA mutation with progerin-like protein accumulation OR ii) Homozygous or compound heterozygous ZMPSTE24 mutations. Patient has a body surface area of 0.39 m <sup>2</sup> and above.
Age Restrictions	Patient is 12 months of age or older.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

## ZOLADEX (S)

### Products Affected

- Zoladex INJ 3.6MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prostate Cancer: Diagnosis of advanced or metastatic prostate cancer. Trial and failure, contraindication, or intolerance to any brand Lupron formulation. Endometriosis [Zoladex (3.6 mg strength)]: Treatment of endometriosis. Trial and failure to Lupron Depot (3.75 mg or 11.25 mg). Advanced Breast Cancer [Zoladex (3.6 mg strength)]: For the palliative treatment of advanced breast cancer. Endometrial thinning [Zoladex (3.6 mg strength)] For the treatment of dysfunctional uterine bleeding. Used as an endometrial thinning agent prior to endometrial ablation.
Age Restrictions	Endometriosis: 18 years or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## ZOLINZA (S)

---

### Products Affected

- Zolinza

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cutaneous T-cell lymphoma (CTCL): Diagnosis of CTCL. Progressive, persistent or recurrent disease on or contraindication or intolerance to two systemic therapies (e.g., bexarotene, romidepsin, etc.).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist/oncologist.
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

# ZYDELIG (S)

## Products Affected

- Zydelig

PA Criteria	Criteria Details
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic lymphocytic leukemia (CLL): Diagnosis of CLL. Used in combination with Rituxan (rituximab). The patient has relapsed on at least one prior therapy (eg, purine analogues [fludarabine, pentostatin, cladribine], alkylating agents [chlorambucil, cyclophosphamide], or monoclonal antibodies [rituximab]). Patient is a candidate for Rituxan (rituximab) monotherapy due to presence of other comorbidities (eg, coronary artery disease, peripheral vascular disease, diabetes mellitus, pulmonary disease [COPD]). Follicular Lymphoma (FL): Diagnosis of FL. The patient has relapsed on at least two prior systemic therapies (eg, rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicine], purine analogs [fludarabine]). Small lymphocytic lymphoma (SLL): Diagnosis of SLL. The patient has relapsed on at least two prior systemic therapies (eg, rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicine], purine analogs [fludarabine]).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	All uses: Prescribed by or in consultation with an oncologist/hematologist.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve for continuation of prior therapy.

## ZYKADIA (S)

---

### Products Affected

- Zykadia

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC that is metastatic or recurrent. Tumor is anaplastic lymphoma kinase (ALK)-positive as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	Approve for continuation of prior therapy.

## ZYTIGA (PREFERRED) (S)

### Products Affected

- Abiraterone Acetate

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Metastatic Castration-Resistant Prostate Cancer (mCRPC): Diagnosis of metastatic castration-resistant (chemical or surgical) or recurrent prostate cancer. Used in combination with prednisone. Metastatic Castration-Sensitive Prostate Cancer (mCSPC): Diagnosis of metastatic high-risk castration-sensitive prostate cancer. Used in combination with prednisone.
Age Restrictions	N/A
Prescriber Restrictions	mCRPC, mCSPC: Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	mCRPC, mCSPC: 12 months
Other Criteria	Approve for continuation of prior therapy

## PART B VERSUS PART D

---

### Products Affected

- Abelcet
- Acetylcysteine SOLN
- Acyclovir Sodium INJ 50MG/ML
- Akynzeo CAPS
- Albuterol Sulfate NEBU 0.083%, 2.5MG/0.5ML
- Ambisome
- Aminosyn II INJ 107.6MEQ/L; 1490MG/100ML; 1527MG/100ML; 1050MG/100ML; 1107MG/100ML; 750MG/100ML; 450MG/100ML; 990MG/100ML; 1500MG/100ML; 1575MG/100ML; 258MG/100ML; 447MG/100ML; 1083MG/100ML; 795MG/100ML; 50MEQ/L; 600MG/100ML; 300MG/100ML; 405MG/100ML; 750MG/100ML
- Amphotericin B INJ
- Amphotericin B Liposome
- Aprepitant CAPS
- Azathioprine TABS
- Budesonide SUSP
- Clinisol Sf 15%
- Cromolyn Sodium NEBU
- Cyclophosphamide CAPS
- Cyclosporine CAPS
- Cyclosporine Modified
- Engerix-b
- Epoprostenol Sodium INJ 0.5MG
- Everolimus TABS 0.25MG, 0.5MG, 0.75MG, 1MG
- Formoterol Fumarate NEBU
- Ganciclovir INJ 500MG, 500MG/10ML
- Gengraf CAPS 100MG, 25MG
- Gengraf SOLN
- Hyperhep B
- Hyperrab
- Imovax Rabies (h.d.c.v.)
- Ipratropium Bromide INHALATION SOLN 0.02%
- Ipratropium Bromide/albuterol Sulfate
- Mycophenolate Mofetil CAPS
- Mycophenolate Mofetil SUSR
- Mycophenolate Mofetil TABS
- Mycophenolic Acid Dr
- Nabi-hb INJ 312UNIT/ML
- Nutrilipid
- Ondansetron Hcl SOLN
- Ondansetron Hydrochloride TABS
- Ondansetron Odt
- Pentamidine Isethionate INHALATION SOLR
- Perforomist
- Plenamine INJ 147.4MEQ/L; 2.17GM/100ML; 1.47GM/100ML; 434MG/100ML; 749MG/100ML; 1.04GM/100ML; 894MG/100ML; 749MG/100ML; 1.04GM/100ML; 1.18GM/100ML; 749MG/100ML; 1.04GM/100ML; 894MG/100ML; 592MG/100ML; 749MG/100ML; 250MG/100ML; 39MG/100ML; 960MG/100ML
- Prehevbrio
- Prograf PACK
- Rabavert
- Recombivax Hb
- Sandimmune SOLN
- Sirolimus SOLN
- Sirolimus TABS
- Tacrolimus CAPS
- Tobramycin NEBU
- Zortress TABS 1MG



## **Details**

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

# INDEX

## A

Abelcet .....	312
Abiraterone Acetate.....	311
Acetylcysteine .....	312
Actemra .....	1
Actemra Actpen.....	1
Actemra Sc (s).....	1
Actimmune.....	3
Actimmune (s).....	3
Acyclovir Sodium .....	312
Adcirca (s).....	4
Adempas.....	5
Adempas (s).....	5
Afinitor.....	6
Afinitor (s).....	6
Afinitor Disperz.....	7
Afinitor Disperz (s) .....	7
Aimovig.....	8
Aimovig (s) .....	8
Akynzeo .....	312
Albuterol Sulfate .....	312
Aldurazyme .....	10
Aldurazyme (s).....	10
Alecensa .....	11
Alecensa (s).....	11
Alosetron Hydrochloride.....	136
Alpha-1 Proteinase Inhibitor, Non-preferred (s). 12	
Alpha-1 Proteinase Inhibitor, Prolastin (s).....	13
Alunbrig .....	14
Alunbrig (s) .....	14
Alyq.....	4
Ambisome .....	312
Ambrisentan .....	130
Aminosyn II.....	312
Amitriptyline Hcl .....	96
Amitriptyline Hydrochloride.....	96
Amnesteem.....	110

Amphotericin B.....	312
Amphotericin B Liposome.....	312
Ampyra (s) .....	15
Androderm .....	251
Aprepitant .....	312
Aralast Np .....	12
Arcalyst .....	16
Arcalyst (s).....	16
Armodafinil.....	164
Auryxia .....	17
Auryxia (s) .....	17
Austedo .....	18
Austedo (s).....	18
Ayvakit.....	19
Ayvakit (s) .....	19
Azathioprine.....	312

## B

Bafiertam.....	20
Bafiertam (s) .....	20
Balversa.....	21
Balversa (s) .....	21
Benlysta.....	22
Benlysta (s) .....	22
Besremi .....	23
Besremi (s).....	23
Betaseron.....	150
Bexarotene .....	240
Bivigam.....	114
Bosulif.....	24
Bosulif (s).....	24
Braftovi .....	25
Braftovi (s) .....	25
Briviact.....	26
Briviact (s) .....	26
Brukina .....	27
Brukina (s).....	27
Budesonide.....	312

**C**

Cablivi .....	28
Cablivi (s).....	28
Cabometyx .....	29
Cabometyx (s) .....	29
Calquence .....	30
Calquence (s).....	30
Caprelsa .....	31
Caprelsa (s).....	31
Cayston.....	32
Cayston (s).....	32
Cerdelga .....	33
Cerdelga (s) .....	33
Chlorzoxazone.....	95
Cholbam .....	34
Cholbam (s).....	34
Ciclodan .....	35
Ciclopirox (s).....	35
Ciclopirox Nail Lacquer.....	35
Cinryze .....	36
Cinryze (s).....	36
Claravis.....	110
Clinisol Sf 15% .....	312
Clovique .....	232
Cometriq.....	37
Cometriq (s).....	37
Copiktra .....	38
Copiktra (s).....	38
Corlanor.....	39
Corlanor (s) .....	39
Cosentyx.....	40
Cosentyx (s).....	40
Cosentyx Sensoready Pen .....	40
Cotellic .....	42
Cotellic (s).....	42
Cromolyn Sodium .....	312
Cuvitru.....	212
Cyclobenzaprine Hydrochloride .....	95
Cyclophosphamide.....	312
Cyclosporine.....	312
Cyclosporine Modified.....	312
Cystaran.....	43
Cystaran (s) .....	43

**D**

D.h.e. 45 (s).....	44
Dalfampridine Er .....	15
Daliresp .....	45
Daliresp (s).....	45
Danyelza .....	46
Danyelza (s) .....	46
Daraprim (s) .....	47
Darzalex Faspro .....	48
Darzalex Faspro (s) .....	48
Daurismo .....	50
Daurismo (s).....	50
Deferasirox.....	51
Deferasirox (s) .....	51
Deferiprone .....	75
Diacomit.....	52
Diacomit (s) .....	52
Diclofenac Sodium.....	180
Dihydroergotamine Mesylate.....	44, 148
Dimethyl Fumarate .....	246
Dimethyl Fumarate Starterpack .....	246
Doxepin Hcl.....	96
Doxepin Hydrochloride .....	96
Dronabinol .....	144
Droxidopa .....	160
Dupixent.....	53
Dupixent (s) .....	53

**E**

Elaprase.....	55
Elaprase (s).....	55
Emgality.....	56
Emgality (s).....	56
Enbrel.....	58
Enbrel (s).....	58
Enbrel Mini .....	58
Enbrel Sureclick.....	58
Engerix-b.....	312
Enspryng .....	59
Enspryng (s) .....	59
Epclusa Preferred (s).....	60
Epidiolex .....	61
Epidiolex (s).....	61

Epoetin Alfa (s) .....	62
Epoprostenol Sodium .....	312
Erivedge .....	64
Erivedge (s) .....	64
Erleada .....	65
Erleada (s) .....	65
Erlotinib Hydrochloride .....	239
Esbriet .....	66
Esbriet (s) .....	66
Eucrisa .....	67
Eucrisa (s) .....	67
Everolimus .....	6, 7, 312
Evrysdi .....	68
Evrysdi (s) .....	68
Exkivity .....	70
Exkivity (s) .....	70

## **F**

Farydak .....	71
Farydak (s) .....	71
Fasenra .....	72
Fasenra (s) .....	72
Fasenra Pen .....	72
Fentanyl (s) .....	74
Fentanyl Citrate Oral Transmucosal .....	74
Ferriprox (s) .....	75
Fintepla .....	76
Fintepla (s) .....	76
Firazyr (s) .....	77
Firmagon .....	78
Firmagon (s) .....	78
Formoterol Fumarate .....	312
Forteo .....	249
Fotivda .....	79
Fotivda (s) .....	79

## **G**

Galafold .....	80
Galafold (s) .....	80
Gamastan .....	81
Gamastan S/d (s) .....	81
Gammagard S/d Iga Less Than 1mcg/ml .....	114
Ganciclovir .....	312
Gavreto .....	82

Gavreto (s) .....	82
Gengraf .....	312
Genotropin .....	87
Genotropin Miniquick .....	87
Gilenya .....	83
Gilenya (s) .....	83
Gilotrif .....	84
Gilotrif (s) .....	84
Glatiramer Acetate .....	85
Glatiramer Acetate (s) .....	85
Gleevec (s) .....	86
Glydo .....	131
Growth Hormone, Preferred (s) .....	87

## **H**

Herzuma .....	90
Herzuma (s) .....	90
Hrm - Antipsychotics .....	91
Hrm - Megestrol Suspension .....	92
Hrm - Megestrol Tablet .....	93
Hrm - Phenobarbital, Pentobarbital .....	94
Hrm - Skeletal Muscle Relaxants .....	95
Hrm - Tca .....	96
Humira .....	97
Humira (s) .....	97
Humira Pediatric Crohns Disease Starter Pack .....	97
Humira Pen .....	97
Humira Pen-cd/uc/hs Starter .....	97
Humira Pen-pediatric Uc Starter Pack .....	97
Humira Pen-ps/uv Starter .....	97
Hyperhep B .....	312
Hyperrab .....	312

## **I**

Ibrance .....	99
Ibrance (s) .....	99
Icatibant Acetate .....	77
Iclusig .....	100
Iclusig (s) .....	100
Icosapent Ethyl .....	275
Idhifa .....	101
Idhifa (s) .....	101
Ilumya .....	102
Ilumya (s) .....	102

Imatinib Mesylate.....	86
Imbruvica .....	103
Imbruvica (s) .....	103
Imovax Rabies (h.d.c.v.) .....	312
Increlex.....	104
Increlex (s).....	104
Inlyta.....	105
Inlyta (s) .....	105
Inqovi .....	106
Inqovi (s) .....	106
Inrebic.....	107
Inrebic (s) .....	107
Intron A .....	108
Intron A (s).....	108
Ipratropium Bromide.....	312
Ipratropium Bromide/albuterol Sulfate .....	312
Iressa.....	109
Iressa (s) .....	109
Isotretinoin .....	110
Isotretinoin (s) .....	110
Istodax (s).....	111
Isturisa .....	112
Isturisa (s).....	112
Itraconazole .....	218
Ivermectin.....	113
Ivermectin (s) .....	113
Ivig (s) .....	114

## **J**

Jakafi .....	116
Jakafi (s) .....	116
Juxtapid .....	117
Juxtapid (s) .....	117

## **K**

Kalydeco.....	118
Kalydeco (s) .....	118
Kanjinti.....	119
Kanjinti (s).....	119
Kanuma .....	120
Kanuma (s) .....	120
Kesimpta.....	121
Kesimpta (s) .....	121
Kisqali .....	122

Kisqali (s).....	122
Kisqali Femara 200 Dose.....	123
Kisqali Femara 400 Dose.....	123
Kisqali Femara 600 Dose.....	123
Kisqali-femara Pack (s).....	123
Korlym .....	124
Korlym (s).....	124
Koselugo .....	125
Koselugo (s) .....	125
Kuvan (s).....	126
Kynmobi .....	127
Kynmobi (s) .....	127
Kynmobi Titration Kit .....	127

## **L**

Lapatinib Ditosylate.....	266
Lemtrada .....	128
Lemtrada (s) .....	128
Lenalidomide .....	198
Lenvima (s) .....	129
Lenvima 10 Mg Daily Dose.....	129
Lenvima 12mg Daily Dose .....	129
Lenvima 14 Mg Daily Dose.....	129
Lenvima 18 Mg Daily Dose.....	129
Lenvima 20 Mg Daily Dose.....	129
Lenvima 24 Mg Daily Dose.....	129
Lenvima 4 Mg Daily Dose.....	129
Lenvima 8 Mg Daily Dose.....	129
Letairis (s) .....	130
Leuprolide Acetate.....	139
Lidocaine.....	131, 132
Lidocaine Hcl.....	131
Lidocaine Hcl Jelly .....	131
Lidocaine Topical (s) .....	131
Lidocaine/prilocaine .....	131
Lidocaine-prilocaine-cream Base .....	131
Lidoderm (s).....	132
Livmarli.....	133
Livmarli (s) .....	133
Lonsurf.....	134
Lonsurf (s).....	134
Lorbrena .....	135
Lorbrena (s).....	135
Lotronex (s).....	136

Lumakras.....	137
Lumakras (s).....	137
Lumizyme.....	138
Lumizyme (s) .....	138
Lupron (s).....	139
Lupron Depot (1-month) .....	140
Lupron Depot (3-month) .....	140
Lupron Depot (4-month) .....	140
Lupron Depot (6-month) .....	140
Lupron Depot (s) .....	140
Lynparza.....	141
Lynparza Tablet (s) .....	141

## **M**

Makena.....	143
Makena (s).....	143
Marinol (s).....	144
Mavyret .....	145
Mavyret (s) .....	145
Megestrol Acetate .....	92, 93
Mekinist.....	146
Mekinist (s) .....	146
Mektovi .....	147
Mektovi (s) .....	147
Miglustat.....	300
Migranal (s) .....	148
Modafinil.....	188
Monjuvi .....	149
Monjuvi (s).....	149
Ms Interferons (preferred) (s).....	150
Mvasi.....	151
Mvasi (s).....	151
Myalept.....	153
Myalept (s) .....	153
Mycophenolate Mofetil.....	312
Mycophenolic Acid Dr.....	312
Myorisan.....	110

## **N**

Nabi-hb.....	312
Naglazyme.....	154
Naglazyme (s) .....	154
Natpara .....	155
Natpara (s).....	155

Nerlynx .....	156
Nerlynx (s) .....	156
Neulasta.....	157
Neulasta (s) .....	157
Neulasta Onpro Kit .....	157
Nexavar .....	158
Nexavar (s).....	158
Ninlaro .....	159
Ninlaro (s) .....	159
Northera (s) .....	160
Noxafil .....	185
Nubeqa .....	161
Nubeqa (s).....	161
Nuedexta .....	162
Nuedexta (s) .....	162
Nuplazid.....	163
Nuplazid (s).....	163
Nutrilipid.....	312
Nuvigil (s).....	164

## **O**

Octreotide Acetate .....	209
Odomzo.....	165
Odomzo (s).....	165
Ofev.....	166
Ofev (s) .....	166
Ondansetron Hcl .....	312
Ondansetron Hydrochloride.....	312
Ondansetron Odt .....	312
Onureg.....	168
Onureg (s) .....	168
Opsumit.....	169
Opsumit (s).....	169
Orencia.....	170, 171
Orencia Clickject .....	171
Orencia IV (s) .....	170
Orencia Sc (s).....	171
Orgovyx .....	172
Orgovyx (s) .....	172
Orkambi .....	173, 174
Orkambi (s) .....	173
Orkambi Granules (s).....	174
Osphena.....	175
Osphena (s) .....	175

Oxandrin (s).....	176
Oxandrolone .....	176
Oxbryta.....	177
Oxbryta (s).....	177

## **P**

Part B Versus Part D .....	312
Pegasys .....	178
Pegasys (s).....	178
Pegasys Proclicl.....	178
Pemazyre .....	179
Pemazyre (s).....	179
Pennsaid (s) .....	180
Pentamidine Isethionate .....	312
Perforomist.....	312
Phenobarbital.....	94
Phesgo .....	181
Phesgo (s) .....	181
Piqray (s) .....	182
Piqray 200mg Daily Dose .....	182
Piqray 250mg Daily Dose .....	182
Piqray 300mg Daily Dose .....	182
Pirfenidone .....	66
Plenamine .....	312
Polivy .....	183
Polivy (s) .....	183
Pomalyst.....	184
Pomalyst (s).....	184
Posaconazole (s).....	185
Posaconazole Dr.....	185
Prehevbrio .....	312
Premium Lidocaine .....	131
Procrit.....	62
Prograf.....	312
Prolastin-c.....	13
Promacta.....	186
Promacta (s).....	186
Provigil (s).....	188
Pyrimethamine .....	47

## **Q**

Qinlock .....	190
Qinlock (s).....	190
Qualaquin (s).....	191

Quinine Sulfate .....	191
-----------------------	-----

## **R**

Rabavert .....	312
Recombivax Hb .....	312
Repatha .....	192
Repatha (s) .....	192
Repatha Pushtronex System.....	192
Repatha Sureclick .....	192
Retevmo .....	195
Retevmo (s).....	195
Revatio (s).....	196
Revcovi .....	197
Revcovi (s) .....	197
Revlimid.....	198
Revlimid (s) .....	198
Rezurock .....	199
Rezurock (s).....	199
Rilutek (s).....	200
Riluzole .....	200
Rinvoq.....	201
Rinvoq (s).....	201
Romidepsin .....	111
Rozlytrek.....	203
Rozlytrek (s).....	203
Rubraca .....	204
Rubraca (s).....	204
Ruxience .....	205
Ruxience (s) .....	205
Rydapt.....	207
Rydapt (s).....	207

## **S**

Sabril (s).....	208
Sajazir .....	77
Sandimmune .....	312
Sandostatin (s).....	209
Sapropterin Dihydrochloride .....	126
Sarclisa.....	210
Sarclisa (s).....	210
Scemblix .....	211
Scemblix (s) .....	211
Scig (s) .....	212
Secuado .....	213

Secuado (s) .....	213
Signifor.....	214
Signifor (s).....	214
Sildenafil Citrate .....	196
Sirolimus .....	312
Skyrizi .....	215
Skyrizi (s) .....	215
Skyrizi Pen .....	215
Sofosbuvir/velpatasvir.....	60
Somatuline Depot.....	216
Somatuline Depot (s).....	216
Somavert.....	217
Somavert (s) .....	217
Sporanox (s) .....	218
Spravato (s) .....	219
Spravato 56mg Dose .....	219
Spravato 84mg Dose .....	219
Sprycel.....	220
Sprycel (s) .....	220
Stelara.....	221, 222
Stelara (iv) (s).....	221
Stelara (s).....	222
Stivarga.....	224
Stivarga (s) .....	224
Strensiq.....	225
Strensiq (s).....	225
Sunitinib Malate .....	227
Supprelin La .....	226
Supprelin La (s).....	226
Sutent.....	227
Sutent (s) .....	227
Sylatron .....	228
Sylatron (s) .....	228
Synagis .....	229
Synagis (s).....	229
Synribo .....	231
Synribo (s).....	231
Syprine (s) .....	232

## **T**

Tabrecta.....	233
Tabrecta (s).....	233
Tacrolimus.....	312
Tadalafil .....	4

Tafamidis (s) .....	234
Tafinlar.....	235
Tafinlar (s) .....	235
Tagrisso.....	237
Tagrisso (s).....	237
Talzenna.....	238
Talzenna (s).....	238
Tarceva (s) .....	239
Targretin.....	240
Targretin (s) .....	240
Tarpeyo .....	241
Tarpeyo (s) .....	241
Tasigna.....	242
Tasigna (s).....	242
Tavalisse .....	243
Tavalisse (s) .....	243
Tavneos .....	244
Tavneos (s).....	244
Tazverik .....	245
Tazverik (s) .....	245
Tecfidera (s) .....	246
Tegsedi.....	247
Tegsedi (s).....	247
Tepmetko .....	248
Tepmetko (s) .....	248
Teriparatide .....	249
Teriparatide (s).....	249
Testosterone .....	251
Testosterone (s).....	251
Testosterone Cypionate.....	251
Testosterone Enanthate .....	253
Testosterone Enanthate (s).....	253
Testosterone Pump.....	251
Tetrabenazine .....	290
Thalomid .....	255
Thalomid (s).....	255
Thioridazine Hcl .....	91
Tibsovo .....	256
Tibsovo (s) .....	256
Tobramycin .....	312
Topical Retinoid (s) .....	257
Trazimera .....	258
Trazimera (s).....	258



Trelstar (s) .....	259
Trelstar Mixject .....	259
Tretinoin .....	257
Trientine Hydrochloride .....	232
Trikafta .....	260
Trikafta (s) .....	260
Triptodur .....	261
Triptodur (s) .....	261
Trodelvy .....	262
Trodelvy (s) .....	262
Truseltiq .....	263
Truseltiq (s) .....	263
Tukysa .....	264
Tukysa (s) .....	264
Turalio .....	265
Turalio (s) .....	265
Tykerb (s) .....	266
Tymlos .....	267
Tymlos (s) .....	267
Tysabri .....	268
Tysabri (s) .....	268

## U

Ubrelvy .....	270
Ubrelvy (s) .....	270
Udenyca .....	271
Udenyca (s) .....	271
Ukoniq .....	272
Ukoniq (s) .....	272

## V

Valchlor .....	273
Valchlor (s) .....	273
Varizig .....	274
Varizig (s) .....	274
Vascepa (s) .....	275
Venclexta .....	276
Venclexta (s) .....	276
Venclexta Starting Pack .....	276
Ventavis .....	277
Ventavis (s) .....	277
Verzenio .....	278
Verzenio (s) .....	278
Vigabatrin .....	208

Vigadrone .....	208
Vimizim .....	279
Vimizim (s) .....	279
Vitrakvi .....	280
Vitrakvi (s) .....	280
Vizimpro .....	281
Vizimpro (s) .....	281
Voriconazole .....	282
Voriconazole Injection (s) .....	282
Vosevi .....	283
Vosevi (s) .....	283
Votrient .....	284
Votrient (s) .....	284
Vyndaqel .....	234

## W

Welireg .....	285
Welireg (s) .....	285

## X

Xalkori .....	286
Xalkori (s) .....	286
Xcopri .....	287
Xcopri (s) .....	287
Xeljanz .....	288
Xeljanz (s) .....	288
Xeljanz Xr .....	288
Xenazine (s) .....	290
Xermelo .....	291
Xermelo (s) .....	291
Xgeva .....	292
Xgeva (s) .....	292
Xifaxan .....	293
Xifaxan (s) .....	293
Xolair .....	294
Xolair (s) .....	294
Xospata .....	296
Xospata (s) .....	296
Xpovio .....	297
Xpovio (s) .....	297
Xpovio 100 Mg Once Weekly .....	297
Xpovio 40 Mg Once Weekly .....	297
Xpovio 40 Mg Twice Weekly .....	297
Xpovio 60 Mg Once Weekly .....	297

Xpovio 60 Mg Twice Weekly .....	297
Xpovio 80 Mg Once Weekly .....	297
Xpovio 80 Mg Twice Weekly .....	297
Xtandi .....	298
Xtandi (s) .....	298
Xyrem .....	299
Xyrem (s) .....	299

## **Z**

Zavesca (s) .....	300
Zejula .....	301
Zejula (s) .....	301
Zelboraf .....	302
Zelboraf (s) .....	302
Zemaira .....	12
Zenatane .....	110

Zepzelca .....	303
Zepzelca (s) .....	303
Zirabev .....	304
Zirabev (s) .....	304
Zokinvy .....	306
Zokinvy (s) .....	306
Zoladex .....	307
Zoladex (s) .....	307
Zolinza .....	308
Zolinza (s) .....	308
Zortress .....	312
Zydelig .....	309
Zydelig (s) .....	309
Zykadia .....	310
Zykadia (s) .....	310
Zytiga (preferred) (s) .....	311